

Boundary conditions and symplectic structure in the Chern-Simons formulation of (2+1)-dimensional gravity

C. Meusburger¹

Perimeter Institute for Theoretical Physics
31 Caroline Street North, Waterloo, Ontario N2L 2Y5, Canada

B. J. Schroers²

Department of Mathematics, Heriot-Watt University
Edinburgh EH14 4AS, United Kingdom

May 2005

Abstract

We propose a description of open universes in the Chern-Simons formulation of (2+1)-dimensional gravity where spatial infinity is implemented as a puncture. At this puncture, additional variables are introduced which lie in the cotangent bundle of the Poincaré group, and coupled minimally to the Chern-Simons gauge field. We apply this description of spatial infinity to open universes of general genus and with an arbitrary number of massive spinning particles. Using results of [9] we give a finite dimensional description of the phase space and determine its symplectic structure. In the special case of a genus zero universe with spinless particles, we compare our result to the symplectic structure computed by Matschull in the metric formulation of (2+1)-dimensional gravity. We comment on the quantisation of the phase space and derive a quantisation condition for the total mass and spin of an open universe.

1 Introduction

The goal of this paper is to introduce a mathematically transparent treatment of spatial infinity for open universes in the Chern-Simons formulation of (2+1)-dimensional gravity [1, 2] with vanishing cosmological constant and to determine the phase space structure of the resulting theory. We consider universes of general genus g and with n massive spinning particles and work with the Hamiltonian formulation of the Chern-Simons formalism.

In general terms, the boundary requirement for an open universe with vanishing cosmological constant which we study in this paper is that there be an asymptotic region of spacetime

¹cmeusburger@perimeterinstitute.ca

²bernd@ma.hw.ac.uk

which has the geometry of a spinning cone [3]. The implementation of this requirement in any Hamiltonian formulation of gravity requires the introduction of boundary degrees of freedom, defined in the asymptotic region of spacetime. These boundary degrees of freedom should then be coupled to the bulk degrees of freedom in such a way that the resulting equations of motion ensure the required conical geometry in the asymptotic region.

The precise formulation of the boundary requirement and its implementation in the metric formulation of gravity in (2+1) dimensions are discussed in detail in [4], which is a key reference for this paper. Technically, the treatment in [4] is quite involved since it is formulated in terms of conditions on the metric in a neighbourhood of spatial infinity. It requires the introduction of infinitely many boundary degrees of freedom, mathematically represented by fields defined in a neighbourhood of spatial infinity. However, after dividing out by gauge degrees of freedom the total phase of the theory turns out to be finite dimensional.

In our treatment of the boundary, the number of degrees of freedom associated to the boundary is finite from the start. This is appropriate for the physics of a spatial boundary in (2+1)-dimensional gravity with vanishing cosmological constant, but should be contrasted with the situation for negative cosmological constant, where one expects infinitely many degrees of freedom to be associated to the boundary of a spacetime containing a black hole [5, 6]. Working in the formulation of (2+1) dimensional gravity as Chern-Simons theory with the universal cover $\tilde{P}_3^\uparrow$ of the Poincaré group as gauge group, we compactify spatial infinity to a point and model it by a puncture on the surface S . More precisely, we consider universes of topology $\mathbb{R} \times S_{g,n}^\infty$, where $S_{g,n}^\infty$ is a oriented surface of genus g with n punctures representing massive, spinning particles. The superscript ∞ refers to the additional puncture which represents the boundary at spatial infinity.

For the first n punctures representing particles we employ the standard treatment of punctures in Chern-Simons theory [7]. We introduce additional variables which lie in coadjoint orbits of the $\tilde{P}_3^\uparrow$, equipped with their standard symplectic structure, and couple them to the gauge field via minimal coupling. Each coadjoint orbit is labelled by the mass and spin of the associated particle. It is explained in [8] why this treatment leads to the correct incorporation of particles in the Chern-Simons formulation of (2+1)-dimensional gravity. An immediate consequence of the minimal coupling to the gauge field is that the curvature of the Chern-Simons gauge field develops a delta-function singularity at each puncture with a coefficient which lies in the coadjoint orbit associated to the puncture.

For the puncture representing spatial infinity, in the following referred to as the distinguished puncture, our treatment differs from the standard approach. We introduce variables which lie in the cotangent bundle of $\tilde{P}_3^\uparrow$, equipped with its canonical symplectic structure, and again couple minimally to the gauge field. Justifying this mathematical description of spatial infinity is one of the main tasks of this paper. In this introduction we only point out that our model leads to a curvature singularity of the Chern-Simons gauge field at the distinguished puncture with a Lie algebra valued coefficient. However, in contrast to the ordinary punctures the Lie algebra valued coefficient is not *a priori* restricted to a fixed adjoint orbit. Thus one

may think of our distinguished puncture as describing an auxiliary particle whose mass and spin are not fixed parameters but additional variables. When all the constraints are imposed, the mass and spin of this auxiliary particle turn out to be minus the total mass and spin of the universe.

Having introduced the distinguished puncture and justified its interpretation as a model for spatial infinity, we parametrise the phase space of the theory and determine its symplectic structure. Here we rely heavily on the paper [9] where, using the method of [10], the phase space structure of Chern-Simons theory with a distinguished puncture is analysed in detail for gauge groups of the form $G \ltimes \mathfrak{g}^*$. We derive some physical consequences of our model and relate our results to other approaches. In particular, we discuss in detail the relation between our description of the phase space and the results derived in the metric formulation of gravity in [4]. In that paper, Matschull considers n spinless particles coupled to gravity. He parametrises the phase space in terms of variables assigned to an oriented graph embedded into a spatial slice of spacetime. We show how to reproduce his result for a particular graph from our description of the phase space in the special case of vanishing genus $g = 0$ and n spinless particles.

We should also comment on the relation between this paper and our investigation of the phase space of open universes in the Chern-Simons formulation of (2+1)-dimensional gravity in [11]. There we considered universes of topology $\mathbb{R} \times (S_{g,n} \setminus D)$, where $S_{g,n} \setminus D$ is a surface of genus g with n punctures and a disc removed. We proposed a parametrisation of the phase space in terms of finitely many holonomy variables and computed the Poisson structure on this parameter space by applying a technique developed by Fock and Rosly [12]. However, the derivation of the Poisson structure given in [11] was not based on a field-theoretical treatment of boundary degrees of freedom and relied partly on symmetry arguments. It turns out that both the symplectic structure associated to the Poisson structure found in [11] and the symplectic structure we compute in this paper can be split into a piece that describes the centre of mass motion of the universe relative to an asymptotic observer and a piece which describes the relative dynamics of the contents of the universe. While the piece describing the relative dynamics is the same in both symplectic structures, the symplectic structures for the centre of mass motion differ. We describe and comment on this difference in Sect. 5.5.

The outline of the paper is as follows. In Sect. 2 we give a brief review of the Chern-Simons formulation of (2+1)-dimensional gravity and introduce our notation and conventions. In Sect. 3 we review in some detail the metric of a spinning cone which surrounds massive point particles with spin in (2+1)-dimensional gravity and describe how this geometry is captured in the Chern-Simons formulation. This section provides important background for Sect. 4, where we review the boundary conditions at spatial infinity, and explain how they lead to the description of spatial infinity as a distinguished puncture. Having formulated our boundary condition for spatial infinity, we then show how this condition can be implemented via an action functional. In Sect. 5 we give a finite-dimensional parametrisation of the phase space and determine the symplectic structure in terms of this parametrisation. The

mathematical results in this section mostly follow directly from our paper [9], but the physical interpretation, particularly in the discussion of gauge fixing, is new. In Sect. 6 we relate our description of the phase to that of Matschull in the special case of vanishing genus $g = 0$ and n spinless particles. Sect. 7 contains remarks about the quantisation of our phase space. In particular, we derive a quantisation condition for the total mass and spin of the universe. Sect. 8 contains our conclusions, and in the appendix we give a brief summary of relevant results from [4] adapted to our conventions.

2 The Chern-Simons formulation of (2+1)-dimensional gravity

2.1 Setting and conventions

We begin with a brief review of the Chern-Simons formulation of (2+1)-dimensional gravity with vanishing cosmological constant for closed universes, referring the reader to [11, 13] for more details. Spacetime is assumed to have the topology $\mathbb{R} \times S_{g,n}$, where $S_{g,n}$ is a two-dimensional, closed and oriented manifold of genus $g \geq 0$ with $n \geq 0$ punctures. We introduce a coordinate x^0 on $\mathbb{R}$ and sometimes we use local coordinates $x = (x^1, x^2)$ on $S_{g,n}$. Differentiation with respect to x^0, x^1 and x^2 is written as ∂_0, ∂_1 and ∂_2 . The coordinates of the n punctures on $S_{g,n}$ are $x_{(1)}, \dots, x_{(n)}$. The symbol d stands for the total exterior derivative of any function or form. Sometimes we consider functions or forms on $\mathbb{R} \times S_{g,n}$ and take exterior derivatives with respect to the dependence on $S_{g,n}$ only; such derivatives are then denoted d_S . Throughout the paper we use units in which the speed of light is 1. In (2+1)-dimensional gravity Newton's constant has dimensions of inverse mass, allowing us to measure masses in units of $(8\pi G)^{-1}$.

In the Chern-Simons formulation, (2+1)-dimensional gravity with vanishing cosmological constant is written as a gauge theory with the (2+1)-dimensional Poincaré group or one of its covers as gauge group. More precisely, writing $L_3^\uparrow$ and $P_3^\uparrow = L_3^\uparrow \ltimes \mathbb{R}^3$ respectively for the proper orthochronous Lorentz and Poincaré group and $\tilde{L}_3^\uparrow, \tilde{P}_3^\uparrow = \tilde{L}_3^\uparrow \ltimes \mathbb{R}^3$ for their universal covers, we are going to take $\tilde{P}_3^\uparrow$ as a gauge group in the following. In order to define the semi-direct product explicitly we need to specify how $\tilde{L}_3^\uparrow$ acts on $\mathbb{R}^3$. We do this by identifying $\mathbb{R}^3$ with the dual of the Lie algebra $so(2, 1)$ of $\tilde{L}_3^\uparrow$. The group $\tilde{P}_3^\uparrow$ is then of the form $G \ltimes \mathfrak{g}^*$ with $G = \tilde{L}_3^\uparrow$ acting on the dual of its Lie algebra in the coadjoint action. This is the formulation we are going to adopt, allowing us to apply the results of the papers [9, 14, 15] where various aspects of Chern-Simons theory with gauge group $G \ltimes \mathfrak{g}^*$ are studied in detail. Since the Lie algebra $so(2, 1)$ is semi-simple, however, we can simplify notation by using the Killing form η on $so(2, 1)$ to identify it with its dual. Explicitly, we introduce generators J_0, J_1, J_2 of $so(2, 1)$ normalised so that

$$\eta(J_a, J_b) = \eta_{ab} = \text{diag}(1, -1, -1) \quad (2.1)$$

and satisfying the commutation relations

$$[J_a, J_b] = \epsilon_{abc} J^c, \quad (2.2)$$

where we use the convention $\epsilon_{012} = 1$ and raised indices with $\eta^{ab} = \eta_{ab}$. To obtain a set of generators for the Lie algebra $\text{Lie } \tilde{P}_3^\uparrow = \text{iso}(2, 1)$ we introduce additional generators P_a , $a = 0, 1, 2$, which commute with each other and transform in the adjoint representation under the generators J_a $a = 0, 1, 2$. The complete commutation relations are then given by

$$[P_a, P_b] = 0, \quad [J_a, J_b] = \epsilon_{abc} J^c, \quad [J_a, P_b] = \epsilon_{abc} P^c. \quad (2.3)$$

The elements P^0, P^1, P^a generate translations in time and space while J_0 generates rotations and J_1 and J_2 Lorentz boosts. The generators J_0 and P^0 of spatial rotations and time translations form an abelian Cartan subalgebra of $\text{iso}(2, 1)$ which we denote by $\mathfrak{c}$ in the following. Note that, because of the Lorentzian signature,

$$[J_0, P_1] = -P_2 \quad \text{and} \quad [J_0, P_2] = P_1, \quad (2.4)$$

so that J_0 generates rotations in the mathematically negative sense in the space spanned by the translation generators P_1 and P_2 .

In this paper only the Cartan subalgebra $\mathfrak{c}$ plays a role, but we should point out that there are non-conjugate Cartan subalgebras in $\text{iso}(2, 1)$, and that some elements in $\text{iso}(2, 1)$ cannot be conjugated into any Cartan subalgebra. The general situation for the Lie algebras of groups of the form $G \ltimes \mathfrak{g}^*$ is summarised in [9], where we also list references. Adopting the terminology used for conjugacy classes of $SO(2, 1)$, we call elements of $\text{iso}(2, 1)$ which are conjugate to $\mu J_0 + s P_0$ for $\mu, s \in \mathbb{R}, \mu \neq 0$, elliptic elements and the Cartan subalgebras conjugate to $\mathfrak{c}$ elliptic Cartan subalgebras. A second family of Cartan subalgebras, called hyperbolic Cartan subalgebras of $\text{iso}(2, 1)$, is obtained from the Cartan subalgebra spanned by J_1 and P_1 by conjugation. Hyperbolic and elliptic Cartan subalgebras cannot be conjugated into each other. Finally, there are elements of the form $h \cdot (\alpha(J_0 \pm J_1) + \beta(P_0 \pm P_1)) \cdot h^{-1}$, for $\alpha, \beta \in \mathbb{R}, h \in \tilde{P}_3^\uparrow$ which are not in any Cartan subalgebra [9, 16]. We call these elements of $\text{iso}(2, 1)$ parabolic.

For the definition of the Chern-Simons action we require a non-degenerate, invariant bilinear form on the Lie algebra. For the Lie algebra $\text{iso}(2, 1)$ we have the pairing

$$\langle J_a, P^b \rangle = \delta_a^b, \quad \langle J_a, J_b \rangle = \langle P^a, P^b \rangle = 0. \quad (2.5)$$

Note that this pairing is canonically defined on the Lie algebra of any Lie group of the form $G \ltimes \mathfrak{g}^*$. It is worth stressing that the earlier identification of $\text{so}(2, 1)$ with $\text{so}(2, 1)^*$ via η is only used to formulate the theory in a more familiar form; the Chern-Simons formulation of gravity requires the bilinear form $\langle, \rangle$, but not the Killing form η .

With our conventions and the parametrisation

$$(u, \mathbf{a}) \in \tilde{P}_3^\uparrow \quad \text{with} \quad u \in \tilde{L}_3^\uparrow, \quad \mathbf{a} \in \mathbb{R}^3, \quad (2.6)$$

the group multiplication law in $\tilde{P}_3^\uparrow$ takes the form

$$(u_1, \mathbf{a}_1) \cdot (u_2, \mathbf{a}_2) = (u_1 \cdot u_2, \mathbf{a}_1 + \text{Ad}(u_1)\mathbf{a}_2) \quad (2.7)$$

with $\text{Ad}(u)$ denoting the $L_3^\uparrow$ element associated to $u \in \tilde{L}_3^\uparrow$. The $\text{Ad}(u)$ action on $\mathbf{a}$ is the familiar representation of an $\tilde{L}_3^\uparrow$ element as an $SO(2, 1)$ matrix. In this paper we will often need to refer to the abelian subgroup of $\tilde{P}_3^\uparrow$ generated by the Cartan subalgebra $\mathfrak{c}$; we will denote it by T_c . Its elements are rotations and time translations and can be parametrised as $(e^{-\Phi J_0}, TP_0)$ with $\Phi, T \in \mathbb{R}$, where the minus sign is inserted to take into account the remark after (2.4).

2.2 The Chern-Simons action

Einstein's theory of gravity is a field theory for a metric g on the spacetime manifold M . The starting point for the Chern-Simons formulation of (2+1)-dimensional gravity is Cartan's point of view, where the theory is formulated in terms of the (non-degenerate) dreibein of one-forms e_a , $a = 0, 1, 2$, and the spin connection one-forms ω_a , $a = 0, 1, 2$. The dreibein is related to the metric via

$$\eta^{ab} e_a \otimes e_b = g, \quad (2.8)$$

and the one-forms ω_a are the coefficients of the spin connection in the expansion

$$\omega = \omega^a J_a. \quad (2.9)$$

In the formulation of (2+1)-dimensional gravity as a Chern-Simons gauge theory, dreibein and spin connection are combined into the Cartan connection [20] or Chern-Simons gauge field. This is a one-form with values in the Lie algebra $iso(2, 1)$

$$A = \omega^a J_a + e_a P^a, \quad (2.10)$$

whose curvature

$$F = T_a P^a + F_\omega^a J_a \quad (2.11)$$

combines the curvature and the torsion of the spin connection:

$$F_\omega = \omega + \frac{1}{2}[\omega, \omega] = F_\omega^a J_a, \quad F_\omega^a = d\omega^a + \frac{1}{2}\epsilon_{bc}^a \omega^b \wedge \omega^c \quad (2.12)$$

$$T_a = de_a + \epsilon_{abc} \omega^b e^c. \quad (2.13)$$

The equations of motion are obtained by variation of an action functional, the Chern-Simons action, which in the absence of punctures (i.e. $n = 0$) is given by

$$S_{CS}[A] = \frac{1}{2} \int_M \langle A \wedge dA \rangle + \frac{2}{3} \langle A \wedge A \wedge A \rangle. \quad (2.14)$$

It is invariant under Chern-Simons gauge transformations

$$A \mapsto \gamma A \gamma^{-1} + \gamma d\gamma^{-1} \quad (2.15)$$

for general functions $\gamma : M \rightarrow \tilde{P}_3^\uparrow$, which take the place of diffeomorphisms in Einstein's formulation of the theory [2]. Variation of the Chern-Simons action (2.14) with respect to the gauge field A yields the condition

$$F = 0. \quad (2.16)$$

Thus, in the absence of matter both the spin connection F_ω and the torsion T vanish. In 2+1 dimensions the vanishing of the spin connection is equivalent to the vanishing of the Einstein tensor. Equation (2.16) is therefore equivalent to the Einstein equations. However, due to the different role of degenerate dreibeins, the precise relationship between Einstein's theory and the Chern-Simons formulation is problematic. In the Cartan formulation of Einstein's theory in 2+1 dimensions, the dreibein e_a is required to define a metric and therefore has to be non-degenerate. The Chern-Simons formulation of (2+1)-dimensional gravity uses the $\tilde{P}_3^\uparrow$ -connection A on M as the basic field and does not impose the non-degeneracy of the dreibein. It is argued in [17], see also [18] for an earlier work in the (1+1)-dimensional context, that this leads to global differences in the structure of the phase spaces of the two theories. The results in this paper are based on the Chern-Simons formulation of (2+1)-dimensional gravity, although we will discuss the link with the metric formulations extensively in Sect. 6.

3 Including particles

3.1 Point particles and the metric of the spinning cone

In order to motivate the coupling of particles to the Chern-Simons gauge field A we need to discuss the metric which solves the Einstein equations for the energy momentum tensor for a single massive particle with spin. We do this in some detail here since this discussion will provide important background for the formulation of boundary conditions in our treatment of open universes in the next section.

The metric which solves Einstein's equations in the presence of a single point particle of mass μ and spin s was derived by Deser, Jackiw and t'Hooft [3]. It is defined on $\mathbb{R}^3$ with the x^0 -axis removed, denoted $\mathbb{R}^3 \setminus \mathbb{R}$ in the following. In terms of polar coordinates r, φ for $x \in \mathbb{R}^2$ so that φ has the range $[0, 2\pi)$, and a third coordinate t the line element can be written

$$ds^2 = (dt + \frac{s}{2\pi}d\varphi)^2 - \frac{1}{(1 - \frac{\mu}{2\pi})^2}dr^2 - r^2d\varphi^2. \quad (3.1)$$

In this solution it is assumed that $0 \leq \mu < 2\pi$. The metric (3.1) is flat everywhere except on the x^0 axis. The quickest way to understand the geometry defined by this metric is to introduce "flat coordinates" T, R, Φ via

$$\Phi = (1 - \frac{\mu}{2\pi})\varphi, \quad T = t + \frac{s}{2\pi}\varphi, \quad R = \frac{r}{1 - \frac{\mu}{2\pi}}, \quad (3.2)$$

in terms of which the line element takes the form of the Minkowski metric

$$ds^2 = dT^2 - dR^2 - R^2d\Phi^2. \quad (3.3)$$

However, the identification $\varphi \sim \varphi + 2\pi$ implies that one has to identify

$$\Phi \sim \Phi + (2\pi - \mu), \quad T \sim T + s. \quad (3.4)$$

If we set $s = 0$ we see that spatial surfaces $T = \text{const.}$ are cones with deficit angle μ and apex on the line defined by $R = 0$. In the general case $s \neq 0$ the geometry defined by (3.3) and (3.4) is called a “spinning cone” with deficit angle μ and time-offset s .

There is a systematic way of introducing the coordinates T, R, Φ discussed under the heading “developing map” in the mathematical literature [19, 20], but also explained in [4] for the context of (2+1)-dimensional gravity. We start with the dreibein and spin connection corresponding to the metric (3.1) given in [8]

$$\begin{aligned} e^0 &= dt + \frac{s}{2\pi} d\varphi & \omega^0 &= \frac{\mu}{2\pi} d\varphi \\ e^1 &= \frac{1}{1-\frac{\mu}{2\pi}} \cos \varphi dr - r \sin \varphi d\varphi & \omega^1 &= 0 \\ e^2 &= \frac{1}{1-\frac{\mu}{2\pi}} \sin \varphi dr + r \cos \varphi d\varphi & \omega^2 &= 0 \end{aligned} \quad \mu \in [0, 2\pi), s \in \mathbb{R}. \quad (3.5)$$

Away from the line $R = 0$ the corresponding $\tilde{P}_3^\dagger$ -gauge field

$$A_p = e_a P^a + \omega^a J_a \quad (3.6)$$

is flat and can therefore be trivialised locally. Since $\mathbb{R}^3 \setminus \mathbb{R}$ is not simply-connected it is not possible to trivialise A_p everywhere in terms of a single $\tilde{P}_3^\dagger$ -valued function. However, defining the multi-valued function

$$\Gamma_p = \left(e^{-\frac{\mu\varphi}{2\pi} J_0}, -(t + \frac{s\varphi}{2\pi}) P_0 - \frac{r}{1-\frac{\mu}{2\pi}} \cos \varphi P_1 - \frac{r}{1-\frac{\mu}{2\pi}} \sin \varphi P_2 \right) \quad (3.7)$$

we have

$$A_p = \Gamma_p d\Gamma_p^{-1}. \quad (3.8)$$

The translational part of the trivialising function defines an embedding of $\mathbb{R}^3 \setminus \mathbb{R}$ into Minkowski space $\mathbb{M}^3$ such that the induced metric is (3.1). Its Lorentz component defines a Lorentz frame at every point of the embedding. Explicitly, parametrising $\Gamma_p^{-1} = (V, \mathbf{X})$ with $V : \mathbb{R}^3 \setminus \mathbb{R} \rightarrow \tilde{L}_3^\dagger$, $\mathbf{X} : \mathbb{R}^3 \setminus \mathbb{R} \rightarrow \mathbb{M}^3$ we have

$$V = e^{\frac{\mu\varphi}{2\pi} J_0} \quad \mathbf{X} = T P_0 + R \cos \Phi P_1 + R \sin \Phi P_2 \quad (3.9)$$

in terms of the coordinates T, R, Φ defined in (3.2). It then follows from (3.8) that $e^a P_a = \text{Ad}(V^{-1}) d\mathbf{X}$ and hence

$$ds^2 = \eta_{ab} e^a e^b = \eta_{ab} dX^a dX^b \quad (3.10)$$

in agreement with (3.3). The embedding map $X : \mathbb{R}^3 \setminus \mathbb{R} \rightarrow \mathbb{M}^3$ shows that one can construct the spinning cone from Minkowski space $\mathbb{M}^3$ by cutting out the wedge $2\pi - \mu < \Phi < 2\pi$ and

identifying the half-planes $\Phi = 2\pi - \mu$ and $\Phi = 0$ after shifting one by s in the T -direction. Equivalently, the spinning cone can be thought of as a quotient of the universal cover of $\mathbb{M}^3 \setminus \mathbb{R}$ by the equivalence relation (3.4).

The function Γ_p in (3.8) is unique only up to right-multiplication with a constant Poincaré element $w^{-1} \in \tilde{P}_3^\uparrow$ or, equivalently, left-multiplication of Γ_p^{-1} with w :

$$\Gamma_p^{-1} \mapsto w\Gamma_p^{-1}, \quad (3.11)$$

which leaves the gauge field (3.8) invariant. Such a transformation corresponds to a Poincaré transformation in the embedding Minkowski space. It leaves the dreibein and spin connection, and hence the metric invariant, but changes the way the cone is embedded. In particular, transformations with elements $w = (e^{-\Phi_0 J_0}, T_0 P_0)$ in the abelian subgroup T_c map $\Phi \mapsto \Phi + \Phi_0$ and $T \mapsto T + T_0$. Such transformations rotate and translate the embedded cone into itself, but more general transformations $w \in \tilde{P}_3^\uparrow$ translate and boost the cone's axis in the Minkowski space. Note that in order to respect the identification (3.4) the $\tilde{P}_3^\uparrow$ -element w itself should be viewed as an element of the quotient of $\tilde{P}_3^\uparrow$ by the relation

$$w \sim w(e^{-(2\pi-\mu)J_0}, sP_0). \quad (3.12)$$

3.2 The spinning cone in the gauge theoretical formulation

In $\tilde{P}_3^\uparrow$ -Chern-Simons theory the gauge field A_p (3.6) should be identified with any field obtained from it via a gauge transformation (2.15). In particular, A_p is gauge equivalent to the singular gauge field

$$\hat{A}_p = \left(\frac{\mu}{2\pi}J_0 + \frac{s}{2\pi}P_0\right)d\varphi \quad (3.13)$$

to which it is related via the gauge transformation

$$A_p = \gamma_p \hat{A}_p \gamma_p^{-1} + \gamma_p d\gamma_p^{-1} \quad (3.14)$$

with the (single-valued) function

$$\gamma_p = \left(1, -tP_0 - \frac{r}{1-\frac{\mu}{2\pi}} \cos \varphi P_1 - \frac{r}{1-\frac{\mu}{2\pi}} \sin \varphi P_2\right). \quad (3.15)$$

This allows us to decompose the trivialising function $\Gamma_p = \gamma_p \hat{\Gamma}_p$ into a single-valued part (3.15) and a multi-valued part

$$\hat{\Gamma}_p = \left(e^{-\frac{\mu\varphi}{2\pi}J_0}, -\frac{s\varphi}{2\pi}P_0\right) \quad (3.16)$$

which trivialises the singular gauge field (3.13)

$$\hat{A}_p = \hat{\Gamma}_p d\hat{\Gamma}_p^{-1}. \quad (3.17)$$

As the translational part of $\hat{\Gamma}_p^{-1}$ is simply $\frac{s\varphi}{2\pi}P_0$, the function $\hat{\Gamma}_p^{-1}$ embeds all points on the cone into the singularity $R = 0$. Unlike (3.9), this map is degenerate and does not induce a

metric on $\mathbb{R}^3 \setminus \mathbb{R}$. The singular gauge illustrates that the metric interpretation of a connection A_p is gauge dependent in $\tilde{P}_3^\dagger$ -gauge theory.

The natural gauge-covariant object associated to a connection is its curvature. For the singular gauge field $\hat{A}_p$ we find

$$\hat{F}_p(x) = (\mu J_0 + sP_0) dx^1 \wedge dx^2 \delta^{(2)}(x). \quad (3.18)$$

Under gauge transformations (2.15) the curvature is conjugated, but since

$$\gamma_p(r=0) = (1, -tP_0) \quad (3.19)$$

the formula (3.18) also gives the curvature of A_p . Under a general gauge transformation γ , the curvature $\hat{F}_p$ at the singularity changes to

$$F_p = h(\mu J_0 + sP_0) h^{-1} dx^1 \wedge dx^2 \delta^{(2)}(x) \quad h(x^0) = \gamma(x^0, 0). \quad (3.20)$$

This formula shows that the gauge invariant object associated to a particle with mass μ and spin s is the coadjoint orbit of the element $\mu J_0 + sP_0$ in the Cartan subalgebra $\mathfrak{c}$, identified with a corresponding adjoint orbit via the non-degenerate form $\langle, \rangle$. Defining

$$T = k_a P^a + p^a J_a = h(\mu J_0 + sP_0) h^{-1} \quad (3.21)$$

and parametrising $h = (v, \mathbf{x})$ we have the explicit formulae

$$p^a J_a = \mu \text{Ad}(v) J_0 \quad k_a P^a = [\mathbf{x}, p^a J_a] + s \text{Ad}(v) P_0, \quad (3.22)$$

or, in terms of the component vectors $\vec{k} = (k_0, k_1, k_2)$, $\vec{x} = (x_0, x_1, x_2)$, $\vec{p} = (p_0, p_1, p_2)$ and $\hat{p} = \vec{p}/\mu$

$$\vec{k} = \vec{x} \wedge \vec{p} + s\hat{p}. \quad (3.23)$$

The vectors $\vec{p}$ and $\vec{k}$ have the physical interpretation of the momentum and (generalised) angular momentum of a free relativistic particle, see [8] and [11] for a more detailed discussion. From (3.22) it follows that they satisfy two relations, the mass and spin constraint

$$p_a p^a = \mu^2 \quad p_a k^a = \mu s. \quad (3.24)$$

We have seen that particles lead to curvature singularities in the formulation of (2+1)-dimensional gravity as a $\tilde{P}_3^\dagger$ -gauge theory. The physical information about the particle, its three-momentum $\vec{p}$ and angular momentum $\vec{k}$, are encoded in the $iso(2, 1)$ -valued coefficient of the curvature at the singularity. An alternative way of capturing the physical information about the particle is the holonomy along a path surrounding the particle. Such holonomies will play an important role in this paper, and we therefore review them briefly. The holonomy for an infinitesimal circle surrounding the puncture in the singular gauge (3.13) is

$$\widehat{\text{Hol}}_p = (e^{-\mu J_0}, -sP_0), \quad (3.25)$$

and in the general gauge (3.14)

$$\text{Hol}_p = h(e^{-\mu J_0}, -sP_0)h^{-1}. \quad (3.26)$$

Parametrising the holonomy as

$$\text{Hol}_p = (u, -\text{Ad}(u)\mathbf{j}) \quad (3.27)$$

and setting $h = (v, \mathbf{x})$, we have

$$u = e^{-p^a J_a} \quad \mathbf{j} = (1 - \text{Ad}(u^{-1}))\mathbf{x} + s\text{Ad}(v)P_0 \quad (3.28)$$

with $\mathbf{p}$ defined as in (3.22). The associated component vector $\vec{\mathbf{j}} = (j_0, j_1, j_2)$ is therefore given by

$$\vec{\mathbf{j}} = (1 - \text{Ad}(u^{-1}))\vec{\mathbf{x}} + s\hat{p}, \quad (3.29)$$

and the vectors $\vec{p}, \vec{\mathbf{j}}$ again satisfy the mass and spin constraint

$$p_a p^a = \mu^2 \quad p_a j^a = \mu s. \quad (3.30)$$

The mathematical motivation of the parametrisation (3.27) comes from the Poisson-Lie structure of $\tilde{P}_3^\dagger$ and is explained in [11] and [9]. Physically, u can be viewed as a group valued momentum and $\mathbf{j}$ as a generalised angular momentum of a point particle coupled to (2+1)-dimensional gravity. The relation between u and $\mathbf{j}$ on the one hand and the free particle momentum $\mathbf{p}$ and angular momentum $\mathbf{k}$ is discussed in some detail in [11].

3.3 Coupling particles to the Chern-Simons action

There is a standard prescription for coupling coadjoint orbits to the Chern-Simons action (2.14), valid for any gauge group [7]. The coupling of particles to (2+1)-dimensional gravity in the Chern-Simons formulation given in [8] follows this prescription. The kinetic term for each particle is derived from the symplectic structure canonically associated to the orbit (3.21), and the orbit parameter h is coupled to the gauge field via minimal coupling. We now consider a spacetime $M \approx \mathbb{R} \times S_{g,n}$ with n particles of mass and spin μ_i and s_i , $i = 1, \dots, n$ and parametrise the associated coadjoint orbit as

$$T_i = h_i(\mu_i J_0 + s_i P_0)h_i^{-1}, \quad h_i \in \tilde{P}_3^\dagger. \quad (3.31)$$

The product structure $M \approx \mathbb{R} \times S_{g,n}$ allows one to give a Hamiltonian formulation of Chern-Simons theory coupled to coadjoint orbits. For this, one decomposes the gauge field with respect to the coordinate x^0 as

$$A = A_0 dx^0 + A_S, \quad (3.32)$$

where A_S is an x^0 -dependent and Lie algebra valued one-form on the spatial surface $S_{g,n}$ and A_0 is a Lie algebra valued function on $\mathbb{R} \times S_{g,n}$. Correspondingly, the curvature F is given as the sum

$$F = dx^0 \wedge (\partial_0 A_S - d_S A_0 + [A_0, A_S]) + F_S, \quad (3.33)$$

of a term proportional to dx^0 and the curvature two-form F_S on $S_{g,n}$

$$F_S = d_S A_S + A_S \wedge A_S. \quad (3.34)$$

In terms of this decomposition, the action for the Chern-Simons formulation of (2+1)-dimensional gravity coupled to particles can be written as

$$\begin{aligned} S[A_S, A_0, h_i] &= \int_{\mathbb{R}} dx^0 \int_{S_{g,n}} \frac{1}{2} \langle \partial_0 A_S \wedge A_S \rangle - \int_{\mathbb{R}} dx^0 \sum_{i=1}^n \langle \mu_i J_0 + s_i P_0, h_i^{-1} \partial_0 h_i \rangle \\ &+ \int_{\mathbb{R}} dx^0 \int_{S_{g,n}} \langle A_0, F_S - \sum_{i=1}^n T_i \delta^{(2)}(x - x_{(i)}) dx^1 \wedge dx^2 \rangle. \end{aligned} \quad (3.35)$$

The Lie algebra valued function A_0 plays the role of a Lagrange multiplier and varying it leads to the constraint

$$F_S(x^0, x) = \sum_{i=1}^n T_i(x^0) \delta^{(2)}(x - x_{(i)}) dx^1 \wedge dx^2. \quad (3.36)$$

The evolution equations obtained by varying A_S are

$$\partial_0 A_S = d_S A_0 + [A_S, A_0], \quad (3.37)$$

and variation of h_i implies

$$\partial_0 T_i = [T_i, A_0(x^0, x_{(i)})]. \quad (3.38)$$

4 Spatial infinity as a distinguished puncture

4.1 Physical motivation

In this section we propose a Chern-Simons model for open universes. Our approach is motivated by Matschull's treatment of the boundary in the metric formulation of (2+1)-dimensional gravity in [4]. That paper focuses on spacetime manifolds of topology $\mathbb{R} \times N$, where N is $\mathbb{R}^2$ with n discs removed. The asymptotic region of N represents spatial infinity, and the boundary condition is the requirement that there exists a neighbourhood of infinity $P_\infty \subset N$ such that the dreibein and spin connection restricted to P_∞ are those of a spinning cone (3.5) in suitable coordinates. This condition can be viewed as the (2+1)-dimensional analogue of requiring that a spacetime is asymptotically Minkowski space in (3+1) dimensions. In physical terms, it states that to an asymptotic observer, the universe

appears like a single particle of mass μ and spin s . However, the cone's deficit angle μ and time offset s are now promoted from constants to phase space variables. As explained in [4] and [21] the introduction of μ and s as phase space variables has to be accompanied by the introduction of further variables in order to obtain a symplectic phase space. These extra variables specify the way the cone is embedded in an Minkowski space. Equivalently, they specify the relation between a distinguished reference frame associated to spatial infinity and the reference frame of an asymptotic observer.

Before explaining the preceding statement in detail, it is useful to consider the analogous situation in (3+1) dimensions, discussed in [22]. There, one usually imposes as a boundary condition that the metric asymptotically takes the form of the (3+1)-dimensional Minkowski metric. The variables associated to spatial infinity are a momentum four-vector and a position four-vector. The components of these two vectors are conjugate to each other and specify, respectively, the universe's total energy and momentum and a time and position with respect to a distinguished reference Minkowski frame associated to spatial infinity. In other words, they specify a Poincaré transformation relating the distinguished Minkowski frame defined by the asymptotic Minkowski metric to the Minkowski frame of an asymptotic observer. Concretely, one can think of such a distinguished Minkowski frame as being given by a set of very distant fixed stars with respect to which an observer can specify time and positions.

In (2+1) dimensions with the boundary condition that the metric is asymptotically conical, the geometry of the asymptotic cone selects a family of frames whose time axis coincides with the cone's symmetry axis. To distinguish one of these frames we need to fix an origin of time and a direction in space by referring to some physical event, in analogy to the (3+1)-dimensional situation reviewed above. The variables associated to spatial infinity then specify a (2+1)-dimensional Poincaré transformation with respect to the distinguished reference frame. In the Chern-Simons formulation, this distinguished reference frame is given by the trivialising function Γ_p in (3.8). The map (3.9) uses coordinates T, R, Φ which implicitly refer to a standard reference frame of $\mathbb{M}^3$ with the point $T = R = 0$ as the origin, with the time axis along the T -axis and the X^1 -axis in the direction $\Phi = 0$. The T -axis is uniquely associated to the embedded cone because it is its symmetry axis; the point $T = 0$ on that axis and the orthogonal direction Φ are fixed by mere convention. We can go from the standard frame to another frame with the same T -axis by rotations and time translations

$$\Phi \mapsto \Phi + \Phi_0, \quad T \mapsto T + T_0, \quad (4.1)$$

which correspond to right-multiplication of the trivialising function (3.8) by elements of the abelian subgroup T_c .

More generally, composing the embedding with the Poincaré transformation w as in (3.11) boosts and translates the axis of the embedded cone with respect to the T -axis. For non-trivial boosts, the axis of the embedded cone is no longer parallel to the T -axis and therefore the intersection point of the axis with surfaces of constant T varies with T . The spacetime asymptotically appears like a particle of mass μ and spin s that moves with respect to an

observer at rest in the reference Minkowski frame with coordinates T, R, Φ . The frames where the axis of the cone coincide with the T -axis can therefore be viewed as centre of mass frames of the universe: the spacetime asymptotically appears like particle of mass μ and spin s at rest at the origin. The set of all centre of mass frames is parametrised by a time shift T_0 and an angle Φ_0 . These are the variables which will play the role of conjugate variables to μ and s as in [4]. At this stage the variables T_0 and Φ_0 merely parametrise the set of centre of mass frames relative to an arbitrarily chosen standard frame. The standard frame only acquires a physical significance and becomes distinguished after gauge fixing in Sect. 5.4.

4.2 The boundary condition at spatial infinity

In this section we show how to implement this treatment of the boundary in the Chern-Simons formulation of (2+1)-dimensional gravity, considering the general case of a spacetime of genus g with n massive, spinning particles. We shall argue shortly that the boundary can be modelled by an additional puncture, which is treated differently from the ordinary punctures representing particles. In order to relate our treatment to the that using the metric formalism in [4] we begin by considering a universe of topology $\mathbb{R} \times (S_{g,n} \setminus D)$.

In the Chern-Simons formulation, the equations of motion derived from the action (3.35) are evolution equations for the connection A_S and the orbit variables h_i . Boundary conditions can be imposed on either A_S or its derivative with respect to x_0 . We seek boundary conditions on A_S which, together with the equations of motion, ensure that, in a suitable gauge, solutions A of the equations of motion have the dreibein and spin connection of a spinning cone in some neighbourhood $\mathbb{R} \times P_\infty$ of the boundary of $\mathbb{R} \times (S_{g,n} \setminus D)$ which does not contain any punctures. To make this statement precise, we introduce polar coordinates $r, \bar{\varphi}$ in P_∞ as follows. Let $\bar{\varphi}$ be an angular coordinate such that closed paths parametrised by increasing $\bar{\varphi} \in [0, 2\pi)$ encircle the removed disc D clockwise, i.e. in the mathematically negative sense. Such paths therefore encircle the “rest of the universe” (handles and ordinary punctures) in a mathematically positive sense, see Fig 1. The coordinate r is a function on P_∞ which increases monotonically towards the boundary and takes some large value at the boundary. Then the requirement that a solution of (2.16) has the asymptotic form of a spinning cone means that there exists a gauge such that the gauge field takes the form

$$A_\infty = e_a P^a + \omega^a J_a \quad (4.2)$$

on $\mathbb{R} \times P_\infty$, with the dreibein and spin connection given by

$$\begin{aligned} e^0 &= dt + \frac{s}{2\pi} d\bar{\varphi} & \omega^0 &= \frac{\mu}{2\pi} d\bar{\varphi} \\ e^1 &= \frac{1}{1-\frac{\mu}{2\pi}} \cos \bar{\varphi} dr - r \sin \bar{\varphi} d\bar{\varphi} & \omega^1 &= 0 \\ e^2 &= \frac{1}{1-\frac{\mu}{2\pi}} \sin \bar{\varphi} dr + r \cos \bar{\varphi} d\bar{\varphi} & \omega^2 &= 0, \end{aligned} \quad (4.3)$$

where t is a function of x^0 .

Our boundary condition is to require the existence of a gauge such that the component A_S of the connection A restricted to P_∞ agrees with the restriction of A_∞ to P_∞ , i.e.

$$A_S = A_{\infty,S} \text{ on } P_\infty. \quad (4.4)$$

As in the discussion preceding (3.13) we can gauge transform this potential into the simpler, but singular form

$$\hat{A}_S = (\frac{\mu}{2\pi}J_0 + \frac{s}{2\pi}P_0)d\bar{\varphi}. \quad (4.5)$$

The singular form of the gauge potential (4.5) captures all the gauge invariant information about the connection in the region P_∞ . We can encode this in a particularly simple fashion by shrinking the disc D to a point x_∞ , called the distinguished puncture in the following, and requiring the curvature to have a delta-function singularity there. For this purpose introduce polar coordinates ρ, φ near the puncture with $\varphi = 2\pi - \bar{\varphi}$ so that the path parametrised by increasing φ encircles the disc D in a mathematically positive sense and ρ is such that the worldline of the puncture is given by $\rho = 0$. With

$$\hat{A}_S = -(\frac{\mu}{2\pi}J_0 + \frac{s}{2\pi}P_0)d\varphi \quad (4.6)$$

the curvature now satisfies

$$\hat{F}_S = -(\mu J_0 + sP_0)dx^1 \wedge dx^2 \delta^{(2)}(x - x_\infty) \quad (4.7)$$

in the region P_∞ . Under gauge transformations $\gamma_\infty : P_\infty \rightarrow \tilde{P}_3^\dagger$ in the region P_∞ , the singular gauge potential $\hat{A}_{\infty,S}$ changes to

$$A_{S,\infty} = \gamma_\infty \hat{A}_{S,\infty} \gamma_\infty^{-1} + \gamma_\infty d_S \gamma_\infty^{-1} \quad (4.8)$$

and the curvature $\hat{F}_S$ gets conjugated by $h = \gamma_\infty(x_\infty)$. We therefore conclude that the requirement that space has the asymptotic geometry of a spinning cone can be implemented in gauge-theoretical formulation of (2+1)-dimensional gravity by adding a distinguished puncture at x_∞ representing spatial infinity and demanding that the spatial part of the curvature is

$$F_S = -h(\mu J_0 + sP_0)h^{-1}dx^1 \wedge dx^2 \delta^{(2)}(x - x_\infty) \quad (4.9)$$

for some $h \in \tilde{P}_3^\dagger$, $\mu \in (0, 2\pi)$ and $s \in \mathbb{R}$ and $h \in \tilde{P}_3^\dagger$.

The parameters μ, s are not fixed constants. Instead, they play the role of phase space variables in our model. We will drop the restriction on the range of μ in the following but shall see in the next subsection that for solutions of the equations of motion μ and s are constant. We can therefore recover solutions with the geometrical interpretation of a spinning cone by restricting attention to solutions with $\mu \in (0, 2\pi)$. More generally, however, we only require that the coefficient of the curvature singularity in (4.9) is an elliptic element $T \in iso(2, 1)$. According to our remarks in Sect. 2.1, these can always be parametrised as

$$T = -h(\mu J_0 + sP_0)h^{-1}, \quad (4.10)$$

for some $h \in \tilde{P}_3^\uparrow$ and $\mu, s \in \mathbb{R}$, $\mu \neq 0$.

As anticipated at the beginning of the section, the set of all T of the form (4.10) does not carry a canonical symplectic structure when μ and s are not fixed. We therefore need additional variables associated to spatial infinity which ensure that we obtain a symplectic structure. These variables should parametrise the embedding of the asymptotic cone in Minkowski space. For this, we note that if a gauge field $A = \Gamma_\infty d\Gamma_\infty^{-1}$ is trivialised by a (multi-valued) function $\Gamma_\infty : \mathbb{R} \times P_\infty \rightarrow \tilde{P}_3^\uparrow$, its spatial part is given by $A_S = \Gamma_\infty d_S \Gamma_\infty^{-1}$ and condition (4.4) is invariant under

$$\Gamma_\infty \mapsto \Gamma_\infty \cdot w^{-1}, \quad (4.11)$$

where w is now an arbitrary $\tilde{P}_3^\uparrow$ -valued function of x^0 . We therefore propose to promote w to a dynamical variable and to couple it w to μ and s via the kinetic term

$$- \langle \mu J_0 + s P_0, w^{-1} \partial_0 w \rangle. \quad (4.12)$$

To ensure gauge invariance, we combine this term with the standard kinetic term $\langle \mu J_0 + s P_0, h^{-1} \partial_0 h \rangle$, so that the total kinetic term for the variables h, w, μ, s associated to the distinguished puncture is given by

$$- \langle \mu J_0 + s P_0, w^{-1} \partial_0 w \rangle + \langle \mu J_0 + s P_0, h^{-1} \partial_0 h \rangle. \quad (4.13)$$

Defining a Poincaré element $T \in \tilde{P}_3^\uparrow$ as in (4.10) and setting $g = hw^{-1}$, we can rewrite (4.13) as

$$\langle T, g \partial_0 g^{-1} \rangle, \quad (4.14)$$

which shows in particular that the kinetic term for the variables associated to the puncture does not depend on the way the coefficient of the curvature singularity (4.9) is parametrised. Moreover, the expression (4.14) has a simple geometric interpretation. As explained in [9], this kinetic term is derived from the symplectic potential of the canonical symplectic structure on the cotangent bundle of $\tilde{P}_3^\uparrow$.

4.3 Action and equations of motion

We now couple the kinetic term (4.13) for the distinguished puncture to the Chern-Simons action (3.35) and show that solutions of the resulting equations of motion have the required behaviour near the distinguished puncture representing spatial infinity. We define our action

for spacetimes $M \approx \mathbb{R} \times S_{g,n}^\infty$ as

$$\begin{aligned}
S[A_S, A_0, h_i, \mu, s, h, w] = & \frac{1}{2} \int_{\mathbb{R}} dx^0 \int_{S_{g,n}^\infty} \langle \partial_0 A_S \wedge A_S \rangle \\
& - \int_{\mathbb{R}} dx^0 \sum_{i=1}^n \langle \mu_i J_0 + s_i P_0, h_i^{-1} \partial_0 h_i \rangle + \int_{\mathbb{R}} dx^0 \langle \mu J_0 + s P_0, h^{-1} \partial_0 h \rangle \\
& + \int_{\mathbb{R}} dx^0 \int_{S_{g,n}^\infty} \langle A_0, F_S + h(\mu J_0 + s P_0) h^{-1} \delta^{(2)}(x - x_\infty) dx^1 \wedge dx^2 - \sum_{i=1}^n T_i \delta^{(2)}(x - x_{(i)}) dx^1 \wedge dx^2 \rangle \\
& - \int_{\mathbb{R}} dx^0 \langle \mu J_0 + s P_0, w^{-1} \partial_0 w \rangle.
\end{aligned} \tag{4.15}$$

Varying the action (4.15) with respect to A_0 we obtain the constraint

$$F_S(x) = -h(\mu J_0 + s P_0) h^{-1} \delta^{(2)}(x - x_\infty) + \sum_{i=1}^n T_i \delta^{(2)}(x - x_{(i)}). \tag{4.16}$$

Variation with respect to A_S and h_i yields the equations (3.37) and (3.38). It remains to derive the equations of motion from the variation of the new variables:

$$\delta \mu \Rightarrow \langle h^{-1} A_0(x_0, x_\infty) h + h^{-1} \partial_0 h - w^{-1} \partial_0 w, J_0 \rangle = 0, \tag{4.17}$$

$$\delta s \Rightarrow \langle h^{-1} A_0(x_0, x_\infty) h + h^{-1} \partial_0 h - w^{-1} \partial_0 w, P_0 \rangle = 0, \tag{4.18}$$

$$\delta h \Rightarrow [\mu J_0 + s P_0, h^{-1} A_0(x_0, x_\infty) h + h^{-1} \partial_0 h] = \partial_0 \mu J_0 + \partial_0 s P_0, \tag{4.19}$$

$$\delta w \Rightarrow [\mu J_0 + s P_0, w^{-1} \partial_0 w] = \partial_0 \mu J_0 + \partial_0 s P_0. \tag{4.20}$$

The last equation implies

$$\partial_0 \mu = \partial_0 s = 0 \tag{4.21}$$

$$[w^{-1} \partial_0 w, J_0] = [w^{-1} \partial_0 w, P_0] = 0. \tag{4.22}$$

Mass μ and spin s of the asymptotic cone are therefore constants of motion. Furthermore, combining equations (4.17), (4.18), (4.19), we find that the function A_0 at the puncture is given by

$$A_0(x_0, x_\infty) = h w^{-1} \partial_0 w h^{-1} + h \partial_0 h^{-1} = h w^{-1} \partial_0 (w h^{-1}). \tag{4.23}$$

To see that these equations capture the geometry of a spinning cone in the region $\mathbb{R} \times P_\infty$ we first derive a general formula for gauge fields that solve the equations of motion in $\mathbb{R} \times P_\infty$.

Theorem 4.1 *For any gauge field solving the equations of motion (4.16) to (4.20) there exists a region P_∞ containing the distinguished puncture and with the topology of a punctured disc in which the gauge field can be written as*

$$A = \Gamma d\Gamma^{-1}, \tag{4.24}$$

with a multi-valued trivialising function $\Gamma : \mathbb{R} \times P_\infty \rightarrow \tilde{P}_3^\uparrow$ of the form

$$\Gamma = \gamma_\infty(e^{\frac{\mu\varphi}{2\pi}J_0}, \frac{s\varphi}{2\pi}P_0)w^{-1}, \quad (4.25)$$

where $\gamma_\infty : \mathbb{R} \times P_\infty \rightarrow \tilde{P}_3^\uparrow$, $\gamma_\infty(x_0, x_\infty) = h$, $w : \mathbb{R} \rightarrow \tilde{P}_3^\uparrow$ satisfies (4.22) and μ, s are real constants.

Proof:

Choosing a neighbourhood P_∞ of the distinguished puncture with the topology of a punctured disc, we can trivialise any gauge field as in (4.24) with a multi-valued function $\Gamma : \mathbb{R} \times P_\infty \rightarrow \tilde{P}_3^\uparrow$, which is unique up to right-multiplication with a constant Poincaré element. It is therefore sufficient to show that the function Γ defined in (4.25) yields a solution of the equations of motion near the distinguished puncture. For this, we note that the spatial component of the gauge field $A_S = \Gamma d_S \Gamma^{-1}$ for the trivialising function (4.25) is given by

$$A_S = -\gamma_\infty(\frac{\mu}{2\pi}J_0 + \frac{s}{2\pi}P_0)\gamma_\infty^{-1}d\varphi + \gamma_\infty d\gamma_\infty^{-1}, \quad (4.26)$$

which, as discussed above, is the most general form of a gauge field satisfying the curvature constraint

$$F_S(x_0, x_\infty) = -h(\mu J_0 + sP_0)h^{-1}\delta^{(2)}(x - x_\infty)dx^1 \wedge dx^2. \quad (4.27)$$

The component A_0 near the puncture is given by

$$A_0 = \Gamma \partial_0 \Gamma^{-1} = \gamma_\infty \partial_0 \gamma_\infty^{-1} + \gamma_\infty(e^{\frac{\mu\varphi}{2\pi}J_0}, \frac{s\varphi}{2\pi}P_0)w^{-1}\partial_0 w(e^{-\frac{\mu\varphi}{2\pi}J_0}, \frac{-s\varphi}{2\pi}P_0)\gamma_\infty^{-1}, \quad (4.28)$$

and taking into account (4.22) we find

$$A_0 = \gamma_\infty \partial_0 \gamma_\infty^{-1} + \gamma_\infty w^{-1} \partial_0 w \gamma_\infty^{-1}. \quad (4.29)$$

It follows that equation (3.37) is satisfied and at the distinguished puncture we have (4.23)

$$A_0(x_0, x_\infty) = h\partial_0 h^{-1} + hw^{-1}\partial_0 wh^{-1} = hw^{-1}\partial_0(wh^{-1}), \quad (4.30)$$

which proves the claim. $\square$

The equations of motion derived from the action (4.15) are therefore equivalent to the requirement that the gauge field takes the form (4.24), (4.25) in the region $\mathbb{R} \times P_\infty$. To establish a link with the gauge field (3.5) for the spinning cone, we now consider the x^0 -evolution of w . The general solution of (4.22) is

$$w(x^0) = \hat{w}c(x^0) \quad (4.31)$$

where $\hat{w}$ is a constant element in $\tilde{P}_3^\uparrow$ and c is an x^0 -dependent element of the abelian subgroup T_c . Parametrising c as

$$c = (e^{\frac{\mu\varphi_0}{2\pi}J_0}, (t_0 + \frac{s\varphi_0}{2\pi})P_0), \quad (4.32)$$

with φ_0 and t_0 functions of x^0 , the gauge field (4.24) on $\mathbb{R} \times P_\infty$ takes the form

$$A = \gamma_\infty \left(\frac{\mu}{2\pi} d(\bar{\varphi} + \varphi_0) J_0 + (dt_0 + \frac{s}{2\pi} d(\bar{\varphi} + \varphi_0) P_0) \gamma_\infty^{-1} + \gamma_\infty d\gamma_\infty^{-1} \right), \quad (4.33)$$

where we have again used $\bar{\varphi} = 2\pi - \varphi$. Up to a trivial shift of the angle, (4.33) agrees with (4.8), and we recover the gauge field (3.5) for a spinning cone by setting

$$\gamma_\infty = \left(1, -\frac{r}{1 - \frac{\mu}{2\pi}} \cos(\bar{\varphi} + \varphi_0) P_1 - \frac{r}{1 - \frac{\mu}{2\pi}} \sin(\bar{\varphi} + \varphi_0) P_2 \right), \quad (4.34)$$

where r is a function of ρ only. The function r can be chosen to increase to an arbitrary large value as ρ decreases, but we must have $r = 0$ at the distinguished puncture in order for (4.34) to be well-defined ($\bar{\varphi}$ -independent) there.

We have thus shown that the equations of motion derived from the action (4.15) have solutions which, in a suitably gauge, have the geometrical interpretation of a spinning cone near the distinguished puncture representing spatial infinity. As explained in Sect. 4.1, the Poincaré element $w \in \tilde{P}_3^\uparrow$ specifies the embedding of the cone into Minkowski space. Equations (4.22), (4.31) which restrict the x^0 -evolution of w to right-multiplication with elements of the abelian subgroup T_c therefore imply that the embedding of the cone's symmetry axis does not change. The right-multiplication with x^0 -dependent elements of T_c corresponds to rotations around and translation in the direction of the cone's axis. We will see in Sect. 5.2 that they are related to gauge transformations. In accordance with the remarks made at the end of Sect. 4.1 they only acquire a physical significance after gauge fixing.

The fact that the x^0 -evolution of w is restricted to right-multiplication with elements of T_c allows one in particular to restrict w in the action (4.15) to the abelian subgroup T_c , i. e. to set $\hat{w} = 0$ in (4.31). Geometrically, this amounts to restricting ourselves to embeddings of the cone into Minkowski space such that the axis of the cone coincides with the T -axis, i.e. to centre of mass frames of the universe. With the parametrisation (4.31) and (4.32) we find

$$-\langle \mu J_0 + s P_0, w^{-1} \partial_0 w \rangle = -\mu \partial_0 t_0 - \frac{\mu s}{2\pi} \partial_0 \varphi_0 - \partial_0 \left(\frac{\mu s \varphi_0}{2\pi} \right). \quad (4.35)$$

Inserting this expression into the action (4.15) and dropping the total time derivative, we then obtain the centre of mass action

$$\begin{aligned} S[A_S, A_0, h_i, \mu, s, h, w] &= \frac{1}{2} \int_{\mathbb{R}} dx^0 \int_{S_{g,n}^\infty} \langle \partial_0 A_S \wedge A_S \rangle \\ &- \int_{\mathbb{R}} dx^0 \sum_{i=1}^n \langle \mu_i J_0 + s_i P_0, h_i^{-1} \partial_0 h_i \rangle + \int_{\mathbb{R}} dx^0 \langle \mu J_0 + s P_0, h^{-1} \partial_0 h \rangle \\ &+ \int_{\mathbb{R}} dx^0 \int_{S_{g,n}^\infty} \langle A_0, F_S + h(\mu J_0 + s P_0) h^{-1} \delta^{(2)}(x - x_\infty) dx^1 \wedge dx^2 - \sum_{i=1}^n T_i \delta^{(2)}(x - x_{(i)}) dx^1 \wedge dx^2 \rangle \\ &- \int_{\mathbb{R}} dx^0 (\mu \partial_0 t_0 + \frac{\mu s}{2\pi} \partial_0 \varphi_0). \end{aligned} \quad (4.36)$$

We should stress that φ_0 and t_0 are merely coordinates used for parametrising elements of the abelian group T_c . They have a geometrical interpretation because the gauge field takes

the form (4.33) in terms of them. They will be useful when establishing a link with the metric formulation of gravity in Sect. 6, but are not particularly convenient for discussing gauge invariance and symmetry of our model. For that purpose it is much better to work with the general Poincaré element w , with the identification (3.12).

5 Symplectic structure

5.1 Symplectic structure on the extended phase space

After defining an action functional which implements spatial infinity as a distinguished puncture, we will now investigate a description of the associated symplectic structure on the phase space. The canonical symplectic form corresponding to the action (4.15) is given by

$$\begin{aligned} \Omega = & -\delta\langle\mu J_0 + sP_0, h^{-1}\delta h\rangle + \delta\langle\mu J_0 + sP_0, w^{-1}\delta w\rangle + \sum_{i=1}^n \delta\langle\mu_i J_0 + s_i P_0, h_i^{-1}\delta h_i\rangle \\ & + \frac{1}{2} \int_{S_{g,n}^\infty} \langle\delta A_S \wedge \delta A_S\rangle. \end{aligned} \quad (5.1)$$

It defines a symplectic structure on an auxiliary space, parametrised by the variables $w, h, h_i \in \tilde{P}_3^\uparrow$, the particle's masses μ_i and spins s_i , total mass μ and spin s of the universe and the spatial gauge field A_S . The physical phase space $\mathcal{P}$ is obtained from this auxiliary space by imposing the constraint (4.16) on the curvature of the spatial gauge field A_S , fixing the values of the particle's masses μ_i and spins s_i and dividing by the associated gauge transformations. However, in practice it proves difficult to parametrise the physical phase space explicitly and to express its symplectic structure in terms of a set of independent parameters. We therefore describe its symplectic structure in terms of a two-form Ω on an extended phase space $\mathcal{P}_{ext}$, on which most, but not all gauge freedom is eliminated.

Our description is based on the results in [9], where we applied the method introduced by Alekseev and Malkin [10] to derive such a description of the phase space for the general case of Chern-Simons theory with gauge groups $G \ltimes \mathfrak{g}^*$ on manifolds $M \approx \mathbb{R} \times S_{g,n}^\infty$. The action functional considered in [9] with the choice $G \ltimes \mathfrak{g}^* = \tilde{P}_3^\uparrow$ gives (4.15), and we can therefore directly apply the results from [9]. We then obtain a parametrisation of the phase space in terms of the total mass μ and spin s of the universe, the particle's masses and spins μ_i, s_i and the Poincaré element $w \in \tilde{P}_3^\uparrow$ together with $n + 2g$ $\tilde{P}_3^\uparrow$ -valued variables closely related to the holonomies of a set of generators of the fundamental group $\pi_1(S_{g,n}^\infty)$. These variables, in the following referred to as holonomy variables, are obtained as follows. One chooses an arbitrary base point $p_0 \in S_{g,n}^\infty$ on the spatial surface and considers a set of curves M_i, A_j, B_j , $i = 1, \dots, n$, $j = 1, \dots, g$ based at p_0 whose homotopy equivalence classes generate the fundamental group $\pi_1(S_{g,n}^\infty, p_0)$, see Fig. 1. As their homotopy equivalence classes generate the fundamental group freely, the holonomies A_j, B_j associated to the curves a_j, b_j around the handles are general elements of the group $\tilde{P}_3^\uparrow$, while the holonomies M_i lie in fixed conjugacy classes

$$\mathcal{C}_{\mu_i s_i} = \{g_i(e^{-\mu_i J_0}, -s_i P_0)g_i^{-1} | g_i \in \tilde{P}_3^\uparrow\} \quad (5.2)$$




Figure 1: The generators of the fundamental group $\pi_1(S_{g,n}^\infty, p_0)$

determined by the masses and spins of the particles. The holonomy L_∞ of the curve l_∞ around the distinguished puncture with respect to the base point p_0 is given as a function of the group elements M_i, A_j, B_j by

$$L_\infty^{-1} = [B_g, A_g^{-1}] \cdots [B_1, A_1^{-1}] \cdot M_n \cdots M_1, \quad (5.3)$$

and can be parametrised in terms of the mass μ and spin s of the universe and a general element $g_\infty \in \tilde{P}_3^\dagger$ as

$$L_\infty^{-1} = g_\infty(e^{-\mu J_0}, -sP_0)g_\infty^{-1}. \quad (5.4)$$

Note that the element $g_\infty \in \tilde{P}_3^\dagger$ in the parametrisation (5.4) is not unique and determined only up to right-multiplication with elements of the abelian subgroup T_c

$$g_\infty \mapsto g_\infty(e^{-\psi J_0}, \tau P_0) \quad \Rightarrow \quad L_\infty^{-1} \mapsto L_\infty^{-1}. \quad (5.5)$$

It will become apparent in Sect. 5.2 that such transformations are related to gauge transformations acting on the extended phase space. We now introduce another set of $\tilde{P}_3^\dagger$ -valued variables $\tilde{M}_i, \tilde{A}_j, \tilde{B}_j$ obtained by conjugating the holonomies M_i, A_j, B_j with g_∞^{-1}

$$\tilde{M}_i = g_\infty^{-1} M_i g_\infty \quad \tilde{A}_j = g_\infty^{-1} A_j g_\infty \quad \tilde{B}_j = g_\infty^{-1} B_j g_\infty. \quad (5.6)$$

From this definition and (5.4) it follows that these variables are subject to the relation

$$[\tilde{B}_g, \tilde{A}_g^{-1}] \cdots [\tilde{B}_1, \tilde{A}_1^{-1}] \cdot \tilde{M}_n \cdots \tilde{M}_1 = (e^{-\mu J_0}, -sP_0). \quad (5.7)$$

We parametrise these holonomy variables as in (3.27) in terms of Lorentz transformations $\tilde{v}_{M_i}, \tilde{u}_{A_j}, \tilde{u}_{B_j} \in \tilde{L}_3^\uparrow$ and angular momenta $\tilde{\mathbf{J}}_{M_i}, \tilde{\mathbf{J}}_{A_j}, \tilde{\mathbf{J}}_{B_j} \in \mathbb{R}^3$ according to

$$\begin{aligned}\tilde{M}_i &= (\tilde{u}_{M_i}, -\text{Ad}(\tilde{u}_{M_i})\tilde{\mathbf{J}}_{M_i}) = (\tilde{v}_{M_i}, \tilde{\mathbf{x}}_{M_i})(e^{-\mu_i J_0}, -s_i P_0)(\tilde{v}_{M_i}, \tilde{\mathbf{x}}_{M_i})^{-1} \\ \tilde{A}_i &= (\tilde{u}_{A_i}, -\text{Ad}(\tilde{u}_{A_i})\tilde{\mathbf{J}}_{A_i}) \\ \tilde{B}_i &= (\tilde{u}_{B_i}, -\text{Ad}(\tilde{u}_{B_i})\tilde{\mathbf{J}}_{B_i}),\end{aligned}\tag{5.8}$$

so that for the particles

$$\tilde{u}_{M_i} = \tilde{v}_{M_i} e^{-\mu_i J_0} \tilde{v}_{M_i}^{-1} \quad \tilde{\mathbf{J}}_{M_i} = (1 - \text{Ad}(\tilde{u}_{M_i}^{-1}))\tilde{\mathbf{x}}_{M_i} + \text{Ad}(\tilde{v}_{M_i})s_i P_0.\tag{5.9}$$

The constraint (3.30) then takes the form

$$\langle \text{Ad}(\tilde{v}_{M_i})J_0, \tilde{\mathbf{J}}_{M_i} \rangle - s_i = 0.\tag{5.10}$$

In terms of these variables, our extended phase space $\mathcal{P}_{ext}$ can be characterised as follows

Theorem 5.1 (*Theorem 6.3 in [9]*) *Consider the extended phase space $\mathcal{P}_{ext}$ parametrised by the holonomy variables $\tilde{v}_{M_i}, \tilde{u}_{A_j}, \tilde{u}_{B_j} \in \tilde{L}_3^\uparrow$, $\tilde{\mathbf{J}}_{M_i}, \tilde{\mathbf{J}}_{A_j}, \tilde{\mathbf{J}}_{B_j} \in \mathbb{R}^3$, $w \in \tilde{P}_3^\uparrow$, $\mu, s \in \mathbb{R}$ and the two-form Ω*

$$\Omega = \delta \left(\langle \mu J_0 + s P_0, w^{-1} \delta w \rangle - \frac{1}{2} \mu \delta s + \Theta(\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}, \tilde{u}_{A_j}, \tilde{\mathbf{J}}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{B_j}) \right),\tag{5.11}$$

where Θ is the one-form

$$\begin{aligned}\Theta(\tilde{v}_{M_i}, \tilde{u}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{M_i}, \tilde{\mathbf{J}}_{A_j}, \tilde{\mathbf{J}}_{B_j}) &= \\ &\sum_{i=1}^n \langle \delta(\tilde{u}_{M_{i-1}} \cdots \tilde{u}_{M_1})(\tilde{u}_{M_{i-1}} \cdots \tilde{u}_{M_1})^{-1} - \delta \tilde{v}_{M_i} \tilde{v}_{M_i}^{-1}, \tilde{\mathbf{J}}_{M_i} \rangle \\ &+ \sum_{i=1}^g \langle \delta(\tilde{u}_{H_{i-1}} \cdots \tilde{u}_{H_1} \tilde{u}_{M_n} \cdots \tilde{u}_{M_1})(\tilde{u}_{H_{i-1}} \cdots \tilde{u}_{H_1} \tilde{u}_{M_n} \cdots \tilde{u}_{M_1})^{-1}, \tilde{\mathbf{J}}_{A_i} \rangle \\ &\quad - \langle \delta(\tilde{u}_{A_i}^{-1} \tilde{u}_{B_i}^{-1} \tilde{u}_{A_i} \tilde{u}_{H_{i-1}} \cdots \tilde{u}_{H_1} \tilde{u}_{M_n} \cdots \tilde{u}_{M_1})(\tilde{u}_{A_i}^{-1} \tilde{u}_{B_i}^{-1} \tilde{u}_{A_i} \tilde{u}_{H_{i-1}} \cdots \tilde{u}_{H_1} \tilde{u}_{M_n} \cdots \tilde{u}_{M_1})^{-1}, \tilde{\mathbf{J}}_{A_i} \rangle \\ &+ \sum_{i=1}^g \langle \delta(\tilde{u}_{A_i}^{-1} \tilde{u}_{B_i}^{-1} \tilde{u}_{A_i} \tilde{u}_{H_{i-1}} \cdots \tilde{u}_{H_1} \tilde{u}_{M_n} \cdots \tilde{u}_{M_1})(\tilde{u}_{A_i}^{-1} \tilde{u}_{B_i}^{-1} \tilde{u}_{A_i} \tilde{u}_{H_{i-1}} \cdots \tilde{u}_{H_1} \tilde{u}_{M_n} \cdots \tilde{u}_{M_1})^{-1}, \tilde{\mathbf{J}}_{B_i} \rangle \\ &\quad - \langle \delta(\tilde{u}_{B_i}^{-1} \tilde{u}_{A_i} \tilde{u}_{H_{i-1}} \cdots \tilde{u}_{H_1} \tilde{u}_{M_n} \cdots \tilde{u}_{M_1})(\tilde{u}_{B_i}^{-1} \tilde{u}_{A_i} \tilde{u}_{H_{i-1}} \cdots \tilde{u}_{H_1} \tilde{u}_{M_n} \cdots \tilde{u}_{M_1})^{-1}, \tilde{\mathbf{J}}_{B_i} \rangle\end{aligned}\tag{5.12}$$

with $\tilde{u}_{H_i} = [\tilde{u}_{B_i}, \tilde{u}_{A_i}^{-1}] = \tilde{u}_{B_i} \tilde{u}_{A_i}^{-1} \tilde{u}_{B_i}^{-1} \tilde{u}_{A_i}$. The physical phase space is obtained from $\mathcal{P}_{ext}$ by imposing the constraint (5.7) and the n spin constraints (5.10) and by dividing by the associated gauge transformations. The two-form Ω restricted to the constraint surface is the pull-back of the symplectic structure on the physical phase space to the constraint surface.

We will now discuss the physical interpretation of this result in the context of (2+1)-dimensional gravity. For this, we parametrise the element $w \in \tilde{P}_3^\uparrow$ as

$$w = (v, \mathbf{x})(1, \frac{1}{2}sP_0) = (v, \mathbf{x} + \frac{1}{2}s\text{Ad}(v)P_0) \quad \text{with } v \in \tilde{L}_3^\uparrow, \mathbf{x} \in \mathbb{R}^3,\tag{5.13}$$

so that

$$w^{-1}\delta w = (v^{-1}\delta v, \text{Ad}(v^{-1})\delta \mathbf{x}) + (0, \frac{1}{2}\delta s P_0) + (0, [v^{-1}\delta v, \frac{1}{2}s P_0]). \quad (5.14)$$

The last term gives no contribution to the pairing of $w^{-1}\delta w$ with $\mu J_0 + s P_0$, and the contribution of the second term cancels with the term $-\frac{1}{2}\mu\delta s$ in the two-form (5.11). We can therefore rewrite the two-form (5.11) as the derivative of a one-form

$$\Omega = \delta \left(\langle \mathbf{p}, \delta \mathbf{x} \rangle + \langle s P_0, v^{-1}\delta v \rangle + \Theta(\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}, \tilde{u}_{A_j}, \tilde{\mathbf{J}}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{B_j}) \right). \quad (5.15)$$

with the momentum three-vector

$$\mathbf{p} = \mu \text{Ad}(v) J_0. \quad (5.16)$$

We see that the one-form in (5.11) and (5.15) is the sum of the one-form Θ depending only on the holonomy variables $\tilde{v}_{M_i}, \tilde{u}_{A_j}, \tilde{u}_{B_j} \in \tilde{L}_3^\dagger, \tilde{\mathbf{J}}_{M_i}, \tilde{\mathbf{J}}_{A_j}, \tilde{\mathbf{J}}_{B_j} \in \mathbb{R}^3$ and of terms involving only the variables μ, s and $(v, \mathbf{x}) \in \tilde{P}_3^\dagger$ (equivalently $w \in \tilde{P}_3^\dagger$) associated to spatial infinity. The former describes the relative motion of the different particles and handles in a centre of mass frame of the universe. The terms which depend only on the variables associated to spatial infinity describe the motion of a distinguished centre of mass frame of the universe with respect to an observer (or, equivalently, the motion of an observer relative to a distinguished centre of mass frame of the universe). The geometrical interpretation of the Poincaré transformation $w = (v, \mathbf{x} + \frac{1}{2}s \text{Ad}(v) P_0)$, explained in Sects. 3 and 4.1, is that it specifies the embedding of the spinning cone into Minkowski space. In particular it maps the T -axis to the axis of the embedded cone. Thus the two Poincaré elements w and $(v, \mathbf{x})$ are equivalent in the sense that the associated embeddings differ only by a shift $T \mapsto T + \frac{s}{2}$ along the T -axis.

Our description in terms of the extended phase space with the two-form Ω therefore decouples the relative motion of different particles and handles and the centre of mass motion relative to an observer. The holonomy variables $\tilde{v}_{M_i}, \tilde{u}_{A_j}, \tilde{u}_{B_j} \in \tilde{L}_3^\dagger, \tilde{\mathbf{J}}_{M_i}, \tilde{\mathbf{J}}_{A_j}, \tilde{\mathbf{J}}_{B_j} \in \mathbb{R}^3$ describing the relative motion are linked to the mass μ and spin s of the asymptotic cone by the constraint (5.7). We can view this constraint as the gravitational equivalent of the usual condition specifying the total energy of a system of free particles as a function of the parameters of their relative motion.

5.2 Gauge transformations and symmetries

The physical phase space is obtained from the extended phase space $\mathcal{P}_{ext}$ with variables $\tilde{v}_{M_i}, \tilde{u}_{A_j}, \tilde{u}_{B_j} \in \tilde{L}_3^\dagger, \tilde{\mathbf{J}}_{M_i}, \tilde{\mathbf{J}}_{A_j}, \tilde{\mathbf{J}}_{B_j} \in \mathbb{R}^3, \mu, s \in \mathbb{R}$ and $(v, \mathbf{x}) \in \tilde{P}_3^\dagger$ and two-form (5.15) by imposing the constraints (5.7) and (5.10) and dividing by the associated gauge transformations. We now give explicit expressions for the action of these gauge transformations on the extended phase space.

In a framework where the symplectic structure of the physical phase space is described in terms of a two-form Ω on an extended phase space $\mathcal{P}_{ext}$, gauge invariance manifests itself

in degeneracy of Ω on the constraint surface $\mathcal{C} \subset \mathcal{P}_{ext}$. There exist vector fields Z on the constraint surface, the infinitesimal generators of the gauge transformations, such that the contraction of the two-form Ω with these vector fields is zero on the constraint surface $\iota_Z \Omega = 0$. In the situation at hand, where $\Omega = \delta\chi$ is given as the exterior derivative of a one-form χ as in (5.11), (5.15) and where the gauge transformations are group actions on the extended phase space, there is a rather straightforward way of determining these gauge transformations and their generators directly from the one-form. The method is discussed in detail in [9], Sect. 6, and can be summarised as follows.

Suppose we have the action ρ of a one-parameter group $h(\epsilon)$, $h(0) = 1$, on $\mathcal{C}$, and Z is the vector field generated by $\rho(h(\epsilon))$. Then Z is the infinitesimal generator of a gauge transformation if

$$\iota_Z \delta\chi = 0 \Leftrightarrow \mathcal{L}_Z \chi - \delta\iota_Z \chi = 0. \quad (5.17)$$

To check if this condition holds we consider the change of χ under pull-back with $\rho(h(\epsilon))$, where ϵ is allowed to be a function on $\mathcal{C}$. If the one-form χ changes according to

$$\chi \mapsto \chi + \delta(\epsilon H) + \mathcal{O}(\epsilon^2) \quad (5.18)$$

with a function H on the constraint surface, it follows that $\iota_Z \chi = \chi(Z) = H$ for the associated vector field Z and $\mathcal{L}_Z \chi = \delta H$. Hence, the group action $\rho(h(\epsilon))$ is a gauge transformation if and only if the one-form χ transforms according to (5.18).

We now apply this method to our extended phase space with the two-form (5.15). The first gauge symmetry we encounter is related to the $\tilde{P}_3^\dagger$ -valued constraint (5.7) and arises from the redundancy in parametrising the action (4.15) in terms of the Poincaré elements $h, w \in \tilde{P}_3^\dagger$. It is easy to see that the (4.15) is invariant under simultaneous right-multiplication of w and h by an element of the abelian subgroup T_c , while all other variables stay fixed. On the extended phase space, such a transformation manifests itself in simultaneous right-multiplication of w or, equivalently, $(v, \mathbf{x})$ and conjugation of all holonomy variables

$$\begin{aligned} (v, \mathbf{x}) &\mapsto (v, \mathbf{x})(e^{-\Phi J_0}, TP_0) \\ \tilde{X} &\mapsto (e^{-\Phi J_0}, TP_0)\tilde{X}(e^{\Phi J_0}, -TP_0) \quad \forall \tilde{X} \in \{\tilde{M}_1, \dots, \tilde{B}_g\} \\ \tilde{v}_{M_i} &\mapsto e^{-\Phi J_0} \tilde{v}_{M_i}. \end{aligned} \quad (5.19)$$

This transformation preserves the constraint (5.7) and the spin constraints (5.10) for the particles. As explained in Sect. 4.1, the right multiplication of $(v, \mathbf{x})$ with $(e^{-\Phi J_0}, TP_0)$ corresponds to the rotation around and time translations in the direction of the asymptotic cone's symmetry axis. As anticipated also in Sect. 4.1 such transformations have no physical meaning if we simultaneously rotate and time-translate all other coordinates used in the description of the universe. This is expressed mathematically in the gauge invariance of our symplectic structure under (5.19).

In order to prove that (5.19) is a gauge transformation we first determine the transformation behaviour of the two-form (5.15). Noting that $(v, \mathbf{x}) \mapsto (v, \mathbf{x})(e^{-\Phi J_0}, TP_0)$ is equivalent to

$w \mapsto w(e^{-\Phi J_0}, TP_0)$ we find that under the transformation (5.19)

$$\langle (\mu J_0, sP_0), w^{-1}\delta w \rangle \mapsto \langle (\mu J_0, sP_0), w^{-1}\delta w \rangle + \mu\delta T - s\delta\Phi. \quad (5.20)$$

Furthermore, it follows from Lemma 6.2 in [9], but can also be verified by direct calculation from definitions (5.9) and (5.12), that under (5.19) the one-form (5.12) transforms according to

$$\Theta(\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}, \tilde{u}_{A_j}, \tilde{\mathbf{J}}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{B_j}) \mapsto \Theta(\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}, \tilde{u}_{A_j}, \tilde{\mathbf{J}}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{B_j}) + s\delta\Phi + T\delta\mu. \quad (5.21)$$

The sum $\Theta + \langle (\boldsymbol{\mu}, \mathbf{s}), w^{-1}\delta w \rangle$ therefore changes by the exact one-form $\delta(\mu T)$ and the transformation (5.19) is a gauge transformation.

Another set of gauge transformations is generated by the spin constraints (5.10) for the particles which reflect the redundancy in the parametrisation (5.9) of $\tilde{u}_{M_i}$ in terms of $\tilde{v}_{M_i}$. The elements $\tilde{u}_{M_i}$ are invariant under right-multiplication of the corresponding elements $\tilde{v}_{M_i}$ by a rotation of angle $\Phi_i \in \mathbb{R}$

$$\tilde{v}_{M_i} \mapsto \tilde{v}_{M_i} e^{-\Phi_i J_0} \quad \Phi_i \in \mathbb{R}. \quad (5.22)$$

From formula (5.12), we find that this transformation changes the one-form Θ according to

$$\Theta(\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}, \tilde{u}_{A_j}, \tilde{\mathbf{J}}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{B_j}) \mapsto \Theta(\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}, \tilde{u}_{A_j}, \tilde{\mathbf{J}}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{B_j}) + \sum_{i=1}^n s_i \delta\Phi_i. \quad (5.23)$$

On the constraint surface, where the spins s_i of the particles are all fixed, the last term in (5.23) is a total derivative, and we see that the transformations (5.22) are indeed gauge transformations.

The extended phase space is therefore given by $6(n+2g)$ parameters associated to the holonomy variables in the one-form Θ , $6n$ parameters in the Poincaré element $w \in \tilde{P}_3^\dagger$ together with the mass μ and spin s of the universe. These parameters are subject to six constraints from the condition (5.7) and n spin constraints (5.10) for the particles. Furthermore, there are the two gauge transformations (5.19) as well as n gauge transformations of the form (5.23). The physical phase space is obtained by imposing the constraints and dividing by the gauge transformations. It therefore has dimension

$$6(n+2g) + 6 + 2 - (6+n) - (2+n) = 4n + 12g. \quad (5.24)$$

In particular, for the case $g = 0$, i. e. a system of n particles on a disc, the dimension of the phase space is $4n$, the same as for a system of n free particles without gravitational interaction.

In contrast to the gauge symmetries, physical symmetries are given by group actions $\rho(h(\epsilon))$ on the constraint surface $\mathcal{C}$, which depend only on a *parameter*, not a function ϵ , which leave the two-form Ω invariant and which commute with the gauge symmetries. The last condition ensures that they give rise to group actions on the physical phase space obtained by

taking the quotient with the gauge transformations. (For a detailed discussion of the general relation between symmetries on an extended phase space, gauge symmetries and physical symmetries see [23].) In the case at hand, the physical symmetries are a global action on the (2+1)-dimensional Poincaré group associated to spatial infinity. On the extended phase space, this Poincaré group action is given by

$$(v, \mathbf{x}) \mapsto g(v, \mathbf{x}) \quad g \in \tilde{P}_3^\uparrow \quad (5.25)$$

or, equivalently, $w \mapsto gw$. As the gauge symmetry (5.19) acts on $(v, \mathbf{x})$ and w by right-multiplication and the gauge symmetry (5.22) leaves these variables invariant, we see that the transformation (5.25) indeed commutes with all gauge symmetries. Furthermore, as the parameters in the group action (5.25) are not dependent on the phase space variables $\delta g = 0$, it preserves the term $w^{-1}\delta w$. Using expression (5.11) for the two-form Ω , we find that it leaves Ω invariant and therefore gives rise to a symmetry acting on the physical phase space. This symmetry action reflects the asymptotic Poincaré symmetry of our spacetime. It describes a Poincaré transformation (4.11) acting on the embedding Minkowski space, i. e. a Poincaré transformation with respect to the distinguished reference frame associated to spatial infinity.

5.3 Centre of mass frames

As discussed in Sect. 4.1, the centre of mass frames of the universe, where the axis of the cone coincides with the T -axis in the embedding Minkowski space, are defined by the restriction of the element $w \in \tilde{P}_3^\uparrow$ to the abelian subgroup T_c , i. e. a combination of a rotation and a time translation. Using again the parametrisation (4.32) for such elements, but incorporating a trivial shift in the time coordinate $t_0 \mapsto t_0 - \frac{1}{2}s$, we set

$$(v, \mathbf{x}) = (e^{\frac{\mu\varphi_0}{2\pi}J_0}, (t_0 + \frac{s}{2\pi}\varphi_0)). \quad (5.26)$$

In the centre of mass frames, the momentum $\mathbf{p}$ in (5.15) is directed along the T -axis and, up to a total derivative $\delta(\frac{\mu s}{2\pi}\varphi_0)$, the two-form (5.15) reduces to

$$\Omega = \delta\chi(\tilde{v}_{M_i}, \tilde{u}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{M_i}, \tilde{\mathbf{J}}_{A_j}, \tilde{\mathbf{J}}_{B_j}, \mu, s, t_0, \varphi_0) \quad (5.27)$$

with

$$\chi(\tilde{v}_{M_i}, \tilde{u}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{M_i}, \tilde{\mathbf{J}}_{A_j}, \tilde{\mathbf{J}}_{B_j}, \mu, s, t_0, \varphi_0) = \Theta(\tilde{v}_{M_i}, \tilde{u}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{M_i}, \tilde{\mathbf{J}}_{A_j}, \tilde{\mathbf{J}}_{B_j}) + \mu\delta t_0 + \frac{\mu s}{2\pi}\delta\varphi_0, \quad (5.28)$$

which is the one-form we would have obtained by working directly with action 4.36.

While the gauge transformations (5.22) for each particle take the same form in a centre of mass frame, the gauge transformations (5.19) now correspond to a shift in the time parameter t_0 and the angle φ_0

$$\begin{aligned} t_0 &\mapsto t_0 + \tau & \varphi_0 &\mapsto \varphi_0 + \psi \\ \tilde{X} &\mapsto (e^{\frac{\mu\psi}{2\pi}J_0}, (\tau + \frac{s}{2\pi}\psi)P_0)\tilde{X}(e^{-\frac{\mu\psi}{2\pi}J_0}, -(\tau + \frac{s}{2\pi}\psi)) & \forall \tilde{X} \in \{\tilde{M}_1, \dots, \tilde{B}_g\} \\ \tilde{v}_{M_i} &\mapsto e^{\frac{\mu\psi}{2\pi}J_0}\tilde{v}_{M_i}. \end{aligned} \quad (5.29)$$

Written in terms of φ_0 and t_0 these gauge transformations look more complicated than (5.19), but their interpretation is precisely as explained after (5.19), see also the discussion in the next subsection.

5.4 Gauge fixing and physical degrees of freedom

After deriving an explicit description of the phase space structure in terms of a two-form on the extended phase space, we will now discuss how to eliminate the gauge freedom in our description by imposing appropriate gauge fixing conditions.

We start by considering the gauge transformations (5.22) associated to the spin constraints (5.10) for the particles. These gauge symmetries reflect the redundancy in the parametrisation (5.9), which can in principle be eliminated by requiring that the elements $\tilde{v}_{M_i}$ are pure boosts, parametrised by an angle $\theta_{M_i} \in [0, 2\pi)$ specifying the direction and a boost parameter $\beta_{M_i} \in \mathbb{R}_0^+$

$$\tilde{v}_{M_i} = e^{-\theta_{M_i} J_0} e^{-\beta_{M_i} J_2} e^{\theta_{M_i} J_0}. \quad (5.30)$$

In practice, it is more convenient to work with the redundant parametrisation (5.9) and to keep track of resulting gauge transformations.

We now turn to the gauge transformations (5.19) associated to the constraint (5.7) which relates the holonomy variables to mass μ and spin s of the asymptotic cone. As discussed in Sects. 4.1 and 5.2, these gauge transformations reflect the fact that the centre of mass condition requiring the axis of the asymptotic cone to coincide with the T -axis only determines a coordinate frame up to rotations and time translations. Rotating and time translating both, the holonomy variables describing the relative motion of different particles and handles with respect to an observer and the Poincaré element relating the coordinate frame of the observer to a fixed frame where the axis of the cone coincides with the T -axis as in (5.19) yields an equivalent description of the same physical state. In physical terms, the situation can be described as follows. The observables associated to spatial infinity allow an observer to specify a distinguished reference frame. Once such a frame is specified, motions *with respect to this frame* are physical. However, applying the same transformation to *both* the distinguished frame *and* all other observable objects in the universe does not yield a different state of the universe but a description of the same state in terms of a different coordinate system.

It is useful to again consider the corresponding situation in (3+1)-dimensions. There, one would impose as a boundary condition that the metric is asymptotically the Minkowski metric, and a distinguished frame associated to spatial infinity would usually be defined by referring to a set of very distant fixed stars. The physical information about a system of particles in such a (3+1)-dimensional universe would then be given by its motion relative to these fixed stars. However, applying the same Poincaré transformation to the distinguished frame *and* the motion of the particles would not yield a different state but only a gauge equivalent description of the same state in terms of a different coordinate system. The situation in (2+1)-dimensions is analogous, only that instead of the Poincaré symmetries

associated to Minkowski space, such gauge transformations take values in the symmetry group of the asymptotic cone, the abelian subgroup T_c .

From the discussion above, it is clear that there are two possible ways of gauge fixing. The first is to fix a distinguished reference frame defined by the boundary, i. e. to impose a restriction on the Poincaré element $(v, \mathbf{x}) \in \tilde{P}_3^\uparrow$

$$(v, \mathbf{x}) = (e^{-\theta_\infty J_0} e^{-\beta_\infty J_2} e^{\theta_\infty J_0}, x^1 P_1 + x^2 P_2), \quad \theta_\infty \in [0, 2\pi), \beta_\infty \in \mathbb{R}_0^+. \quad (5.31)$$

A corresponding condition for the centre of mass case would be to set the variables φ_0 and t_0 in (5.28) to zero. However, the gauge fixing condition (5.31) has the disadvantage that is not compatible with the asymptotic Poincaré symmetry (5.25) and yields to complicated expressions for the centre of mass motion.

An alternative gauge fixing condition without these problems is obtained by defining a reference frame in terms of the motion of one of the particles. For instance, assuming that not all particles are at rest relative to the centre of mass of the universe, we can impose without loss of generality that the first particle moves along the 1-axis, which amounts to setting $\theta_{M_1} = 0$ in (5.30)

$$\tilde{v}_{M_1} = e^{-\beta_{M_1} J_2} \quad \beta_{M_1} \in \mathbb{R}_0^+, \quad (5.32)$$

and demand that its angular momentum three-vector is given by

$$\tilde{\mathbf{j}}_{M_1} = \text{Ad}(e^{-\beta_{M_1} J_2}) s_1 P_0 + j_{M_1}^2 P_2, \quad (5.33)$$

which amounts to a restriction on the particle's position vector $\tilde{\mathbf{x}}_{M_1}$. Although in principle it is possible to use the gauge fixing (5.31), (5.32) and (5.33) to express the two-form Ω in terms of independent parameters, there is no obvious way of doing this in practice for a general genus g surface with arbitrarily many massive, spinning particles. For the case of a genus $g = 0$ surface with $n > 1$ massive, but spinless particles, we will see in Sect. 6 that there is a way of solving the constraints which leads to a fairly simple expression for the one-form χ . Spatial surfaces $S_{0,2}^\infty$ with two massive, spinning particles are discussed further in Sect. 7. In this case, the motion of the two particles is determined completely by the mass μ and spin s of the universe and the Poincaré element w .

The centre of mass motion is governed by the symplectic potential

$$\langle (\mu J_0 + s P_0), w^{-1} \delta w \rangle. \quad (5.34)$$

After the imposition of the gauge fixing conditions (5.32), (5.33), the Poincaré element $w \in \tilde{P}_3^\uparrow$ becomes a physically meaningful observable, namely a Poincaré transformation with respect to a fixed frame in which the axis of the cone coincides with the T -axis. As the ambiguity in the definition of w is removed by the gauge fixing conditions (5.32), (5.33), right-multiplication of this element by rotations and time shifts

$$w \mapsto w \cdot (e^{-\Phi J_0}, T P_0) \quad (5.35)$$

now represents rotations and time translations with respect to this reference frame. As discussed in Sect. 3, the angle coordinate Φ and time coordinate T obey the identification

$$(T, \Phi) \sim (T + s, \Phi + 2\pi - \mu) \quad (5.36)$$

leading to the identification (3.12) for the $\tilde{P}_3^\uparrow$ -element w . We will see in Sect. 7 that this condition gives rise to a quantisation condition on mass μ and spin s of the universe.

5.5 Relationship to the flower algebra

To end this section, we briefly explain the relationship between the symplectic structure defined by the two-form (5.11) or, equivalently, (5.15) and the Poisson structure we gave in [11]. As explained in the introduction, that Poisson structure was not derived from a field-theoretical treatment of the boundary. Instead, we argued on symmetry grounds that a Poisson structure defined by Fock and Rosly in [12] as an auxiliary Poisson structure in the study of moduli spaces of flat connections on a surface without boundary could be used to describe the Poisson structure of the phase space of an open universe in the Chern-Simons formulation of (2+1)-dimensional gravity. In [14] we called the Poisson algebra corresponding to that Poisson structure the flower algebra, and we adopt this terminology here. The Poisson structure given in [11] is defined on the space $\left(\tilde{P}_3^\uparrow\right)^{2g+n}$, and it is symplectic when restricted to the space

$$\mathcal{C}_{\mu_1 s_1} \times \dots \times \mathcal{C}_{\mu_n s_n} \times \left(\tilde{P}_3^\uparrow\right)^{2g} \quad (5.37)$$

spanned by the holonomy variables $M_1, \dots, M_n, A_1, B_1, \dots, A_g, B_g$. Parametrising them via

$$\begin{aligned} M_i &= (u_{M_i}, -\text{Ad}(u_{M_i})\mathbf{j}_{M_i}) = (v_{M_i}, \mathbf{x}_{M_i})(e^{-\mu_i J_0}, -s_i P_0)(v_{M_i}, \mathbf{x}_{M_i})^{-1} \\ A_i &= (u_{A_i}, -\text{Ad}(u_{A_i})\mathbf{j}_{A_i}) \\ B_i &= (u_{B_i}, -\text{Ad}(u_{B_i})\mathbf{j}_{B_i}), \end{aligned} \quad (5.38)$$

so that

$$u_{M_i} = v_{M_i} e^{-\mu_i J_0} v_{M_i}^{-1} \quad \mathbf{j}_{M_i} = (1 - \text{Ad}(u_{M_i}^{-1}))\mathbf{x}_{M_i} + \text{Ad}(v_{M_i})s_i P_0, \quad (5.39)$$

the symplectic structure on the symplectic leaf (5.37) is most easily described by pulling it back from the $\tilde{L}_3^\uparrow$ -holonomies $u_{M_1}, \dots, u_{M_n}$ to the $\tilde{L}_3^\uparrow$ -elements $v_{M_1}, \dots, v_{M_n}$. As shown in [15] that pull back is the two-form

$$\Omega_{\text{flower}} = \delta\Theta(v_{M_i}, \mathbf{j}_{M_i}, u_{A_j}, \mathbf{j}_{A_j}, u_{B_j}, \mathbf{j}_{B_j}), \quad (5.40)$$

with Θ defined as in (5.12). To compare Ω_{flower} with Ω in (5.15) we express Ω_{flower} in terms of the holonomies (5.6) and the $\tilde{P}_3^\uparrow$ -element g_∞ . Parametrising $g_\infty = (v_\infty, \mathbf{x}_\infty)$ and

$$g_\infty(e^{-\mu J_0}, -s P_0)g_\infty^{-1} = (u_\infty, -\text{Ad}(u_\infty)\mathbf{j}_\infty) \quad (5.41)$$

we have the equivalent formula

$$\Omega_{\text{flower}} = \delta \left(-\langle \delta u_\infty u_\infty^{-1}, \mathbf{x}_\infty \rangle - \langle s P_0, v_\infty^{-1} \delta v_\infty \rangle + \Theta(\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}, \tilde{u}_{A_j}, \tilde{\mathbf{J}}_{A_j}, \tilde{u}_{B_j}, \tilde{\mathbf{J}}_{B_j}) \right). \quad (5.42)$$

As stated in the introduction, the last term is the same as the term describing the relative motion of handles and particles in the universe in (5.15). The first two terms are analogous to the first two terms in (5.15) describing the centre of mass motion of the universe. However, while our discussion in Sect. 4 has provided a clear interpretation of the Poincaré element w , we have no such interpretation for the Poincaré element g_∞ because the symplectic structure (5.42) was not derived from a field-theoretical treatment of the boundary. We note, however, that we can impose the analogue of the centre of mass condition by requiring g_∞ to take values in the abelian subgroup T_c . Parametrising such as elements as

$$g_\infty = (e^{-\frac{\mu\varphi_0}{2\pi} J_0}, (-t_0 - \frac{s\varphi_0}{2\pi}) P_0), \quad (5.43)$$

we obtain agreement with our symplectic structure (5.27) for centre of mass frames.

6 The link with the metric formulation

Due to the different role of degenerate dreibeins in the two theories, the relation between the phase space of (2+1)-dimensional gravity in its Chern-Simons and its Einstein formulation is rather subtle and not yet fully clarified [17]. In particular, it is not obvious how our description of the phase space for open universes based on the Chern-Simons formalism is related to other approaches which take the metric viewpoint. In this section, we show that our description of the phase space agrees with the results obtained by Matschull [4] who derives a description of the phase space for open universes of genus $g = 0$ with n massive, but spinless particles in a centre of mass frame of the universe.

The derivation in [4] is based entirely on the metric formulation of (2+1)-dimensional gravity and relies on an ADM-decomposition of the spacetime. Spatial infinity is incorporated as a boundary with the boundary condition that dreibein and spin connection asymptotically take the form (3.5) associated to a spinning cone. The formalism makes use of graph embedded into a spatial slice of the spacetime and describes the phase space by means of a symplectic potential on an extended phase space of link variables. For the convenience of the reader, we summarise the results of [4] in the Appendix and show how they can be specialised to a minimal graph.

To establish a link between our description of the phase space and the work [4], we consider manifolds $M \approx \mathbb{R} \times S_{0,n}^\infty$, where the spatial surface is a disc with n massive, but spinless particles and restrict attention to centre of mass frames. In that case, the variables parametrising the extended phase space $\mathcal{P}_{\text{ext}}$ are the total mass μ and spin s of the universe, together with the time and angle coordinates t_0, φ_0 and the holonomy variables $\tilde{v}_{M_1}, \dots, \tilde{v}_{M_n}, \tilde{\mathbf{J}}_{M_1}, \dots, \tilde{\mathbf{J}}_{M_n}$. In terms of these, the symplectic potential on $\mathcal{P}_{\text{ext}}$ in the centre-of-mass frame is given by (5.28), (5.12)

$$\chi(\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}, \mu, s, t_0, \varphi_0) = \Theta(\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}) + \mu \delta t_0 + \frac{\mu s}{2\pi} \delta \varphi_0. \quad (6.1)$$

As discussed in Sect. 5, the variables are subject to a spin constraint (5.10) for each of the particles, where now $s_i = 0$, and the constraint (5.7) with associated gauge transformations (5.22) and (5.19).

It turns out that for the case of n spinless particles on a disc, some of these constraints can be solved explicitly by parametrising the extended phase space in terms of a new set of variables. We have the following lemma, which can be proved by straightforward calculation using the identity $\delta \tilde{u}_{M_i} \tilde{u}_{M_i}^{-1} = \delta \tilde{v}_{M_i} \tilde{v}_{M_i}^{-1} - \tilde{u}_{M_i} \delta \tilde{v}_{M_i} \tilde{v}_{M_i}^{-1} \tilde{u}_{M_i}^{-1}$.

Lemma 6.1

1. For genus $g = 0$ and n spinless particles, the one-form (5.12) takes the form

$$\Theta(\tilde{u}_{M_i}, \tilde{\mathbf{x}}_{M_i}) = \sum_{i=1}^n \langle \delta(\tilde{u}_{M_{i-1}} \cdots \tilde{u}_{M_1})(\tilde{u}_{M_{i-1}} \cdots \tilde{u}_{M_1})^{-1} - \delta(\tilde{u}_{M_i} \cdots \tilde{u}_{M_1})(\tilde{u}_{M_i} \cdots \tilde{u}_{M_1})^{-1}, \tilde{\mathbf{x}}_{M_i} \rangle \quad (6.2)$$

in terms of the variables $\tilde{u}_{M_i}, \tilde{\mathbf{x}}_{M_i}$ related to $\tilde{v}_{M_i}, \tilde{\mathbf{J}}_{M_i}$ by

$$\tilde{u}_{M_i} = \tilde{v}_{M_i} e^{-\mu J_0} \tilde{v}_{M_i}^{-1} \quad \tilde{\mathbf{J}}_{M_i} = (1 - \text{Ad}(\tilde{u}_{M_i}^{-1})) \tilde{\mathbf{x}}_{M_i}. \quad (6.3)$$

2. Introducing a new set of variables via

$$g_1 = \tilde{u}_{M_1}^{-1}, \quad g_i = \tilde{u}_{M_1}^{-1} \cdots \tilde{u}_{M_i}^{-1} \quad i = 2, \dots, n-1 \quad (6.4)$$

$$\mathbf{z}_i = \tilde{\mathbf{x}}_{M_{i+1}} - \tilde{\mathbf{x}}_{M_i} \quad i = 1, \dots, n-1, \quad (6.5)$$

we have the alternative expression

$$\Theta(g_i, \mathbf{z}_i, \tilde{\mathbf{x}}_{M_n}) = - \sum_{i=1}^{n-1} \langle g_i^{-1} \delta g_i, \mathbf{z}_i \rangle - \langle \delta(\tilde{u}_{M_n} g_{n-1}^{-1}) g_{n-1} \tilde{u}_{M_n}^{-1}, \mathbf{x}_{M_n} \rangle. \quad (6.6)$$

In terms of the variables $g_i, \mathbf{z}_i, \tilde{u}_{M_n}$ and $\tilde{\mathbf{x}}_{M_n}$, the constraint (5.7) takes the form

$$\tilde{M}_n \cdots \tilde{M}_1 = (\tilde{u}_{M_n} g_{n-1}^{-1}, -\text{Ad}(\tilde{u}_{M_n} g_{n-1}^{-1}) \tilde{\mathbf{J}}_{K_n}) \approx (e^{-\mu J_0}, -s P_0) \quad (6.7)$$

with

$$\tilde{\mathbf{J}}_{K_n} = (1 - \text{Ad}(\tilde{g}_{n-1} u_{M_n}^{-1})) \tilde{\mathbf{x}}_{M_n} - \sum_{i=1}^{n-1} (1 - \text{Ad}(g_i)) \mathbf{z}_i, \quad (6.8)$$

or, equivalently,

$$\tilde{u}_{M_n} g_{n-1}^{-1} \approx e^{-\mu J_0} \quad (6.9)$$

$$\sum_{i=1}^{n-1} \langle (1 - \text{Ad}(g_i)) \mathbf{z}_i, J_0 \rangle \approx -s \quad (6.10)$$

$$(1 - \text{Ad}(g_{n-1} \tilde{u}_{M_n}^{-1})) \tilde{\mathbf{x}}_{M_n} - \sum_{i=1}^{n-1} (1 - \text{Ad}(g_i)) \mathbf{z}_i + \sum_{i=1}^{n-1} \langle (1 - \text{Ad}(g_i)) \mathbf{z}_i, J_0 \rangle P_0 \approx 0. \quad (6.11)$$

By adding a function of the constraint (6.9), we can rewrite the symplectic potential (6.1) as

$$\chi(g_i, \mathbf{z}_i, \tilde{\mathbf{x}}_{M_n}, \mu, s, t_0, \varphi_0) = - \sum_{i=1}^{n-1} \langle g_i^{-1} \delta g_i, \mathbf{z}_i \rangle + \mu \delta t_0 + \frac{\mu s}{2\pi} \delta \varphi_0 + \langle \tilde{\mathbf{x}}_{M_n}, J_0 \rangle \delta \mu. \quad (6.12)$$

As this expression does not depend on the variable $\tilde{u}_{M_n}$ and only depends on the component of $\tilde{\mathbf{x}}_{M_n}$ parallel to P_0 , we can now solve the constraints (6.9) and (6.11) by setting

$$u_{M_n} = g_{n-1} e^{-\mu J_0} \quad (6.13)$$

$$\tilde{\mathbf{x}}_{M_n} - \langle \tilde{\mathbf{x}}_{M_n}, J_0 \rangle P_0 = \frac{1}{1 - \text{Ad}(e^{\mu J_0})} \sum_{i=1}^{n-1} (1 - \text{Ad}(g_i)) \mathbf{z}_i - \langle (1 - \text{Ad}(g_i)) \mathbf{z}_i, J_0 \rangle P_0, \quad (6.14)$$

where $\frac{1}{1 - \text{Ad}(e^{\mu J_0})}$ denotes the inverse of the bijective map $(1 - \text{Ad}(e^{\mu J_0}))|_{\text{Span}(P_1, P_2)} : \text{Span}(P_1, P_2) \rightarrow \text{Span}(P_1, P_2)$. Furthermore, we can remove the component $\tilde{\mathbf{x}}_{M_n}$ parallel to P_0 by shifting our time parameter

$$T_0 = t_0 - \langle \tilde{\mathbf{x}}_{M_n}, J_0 \rangle. \quad (6.15)$$

This allows one to eliminate the variables $\tilde{u}_{M_n}$, $\tilde{\mathbf{x}}_{M_n}$ from the extended phase space and, after adding a total derivative, to rewrite the symplectic potential (6.12) as

$$\chi(g_i, \mathbf{z}_i, \mu, s, T_0, \varphi_0) = -T_0 \delta \mu + \frac{\mu s}{2\pi} \delta \varphi_0 - \sum_{i=1}^{n-1} \langle g_i^{-1} \delta g_i, \mathbf{z}_i \rangle. \quad (6.16)$$

Our extended phase space is then parametrised by $n-1$ Lorentz transformations g_i , $n-1$ vectors $\mathbf{z}_i \in \mathbb{R}^3$ and the four real parameters μ, s, T_0, φ_0 . While the constraints (6.9) and (6.11) are no longer present, these variables are still subject to the constraint (6.10), and we have traded the spin constraints (5.10) for a mass constraint for each particle. With the definitions $g_n = e^{\mu J_0}$, $g_i^{-1} g_{i-1} = e^{-\tilde{p}_{M_i}^a J_a}$ these mass constraints read

$$\sqrt{\tilde{\mathbf{p}}_{M_i}^2} - \mu_i \approx 0 \quad i = 1, \dots, n. \quad (6.17)$$

Using the methods from Sect. 5.2, we find that the gauge transformations associated to the constraint (6.10) are given by

$$\begin{aligned} \varphi_0 &\mapsto \varphi_0 - \frac{2\pi}{\mu} \psi, & T_0 &\mapsto T_0 + \frac{s}{\mu} \psi \\ g_i &\mapsto e^{-\psi J_0} g_i e^{\psi J_0}, & \mathbf{z}_i &\mapsto \text{Ad}(e^{-\psi J_0}) \mathbf{z}_i, \quad i = 1, \dots, n-1, \end{aligned} \quad (6.18)$$

and the gauge transformations corresponding to the mass constraints (6.17) are

$$\begin{aligned} C_1 : & \quad \mathbf{z}_1 \mapsto \mathbf{z}_1 - \frac{\tau_1}{\mu_1} \tilde{\mathbf{p}}_{M_1} \\ C_i : & \quad \mathbf{z}_i \mapsto \mathbf{z}_i - \frac{\tau_i}{\mu_i} \tilde{\mathbf{p}}_{M_i}, \quad \mathbf{z}_{i-1} \mapsto \mathbf{z}_{i-1} + \frac{\tau_i}{\mu_i} \tilde{\mathbf{p}}_{M_i} \quad i = 2, \dots, n-1 \\ C_n : & \quad \mathbf{z}_{n-1} \mapsto \mathbf{z}_{n-1} + \frac{\tau_n}{\mu_n} \tilde{\mathbf{p}}_{M_n}, \quad T_0 \mapsto T_0 - \frac{\tau_n}{\mu_n} \langle \tilde{p}_{M_n}^a J_a, P^0 \rangle, \end{aligned} \quad (6.19)$$

with parameters $\psi, \tau_i \in \mathbb{R}$. The physical phase space is obtained from the $6(n-1) + 4$ -dimensional extended phase space with symplectic potential (6.16) by imposing the $n+1$ first-class constraints (6.10), (6.17) and dividing by the associated gauge transformations (6.18), (6.19) and therefore has the dimension

$$6(n-1) + 4 - 2(n+1) = 4n - 4. \quad (6.20)$$

This description of the physical phase space in terms of the extended phase space with variables $g_i, z_i, i = 1, \dots, n-1$ and parameters μ, s, T_0, φ_0 and symplectic potential (6.16) is exactly the one obtained by specialising the formalism in [4] to a minimal graph, for details see Sect. A.2 in the appendix. Our description of open universes from the Chern-Simons viewpoint, where spatial infinity is treated as a distinguished puncture therefore agrees with the results in [4] for a system of n spinless particles on a disc in a centre of mass frame of the universe. While the physical interpretation of the results is more readily apparent from the metric viewpoint, our formalism is more general. It gives an efficient description of spacetimes of general genus g with an arbitrary number of particles with non-vanishing spin and without restriction to centre of mass frames. In particular, it is not clear how the formalism in [4] can be extended to include spinning particles, while the inclusion of spin poses no problems in the Chern-Simons framework.

7 The Quantisation condition on mass and spin

After giving a description of the phase space of (2+1)-dimensional gravity for open universes in terms of the two-form Ω on an extended phase space, we will now demonstrate how this description and the results in [14] give rise to a quantisation condition on the total mass μ and spin s of the universe.

It is shown in Sect. 5 that the eight degrees of freedom associated to the centre of mass are the mass μ and spin s of the universe and the Poincaré element $w = (v, \mathbf{x})(1, \frac{1}{2}sP_0) \in \tilde{P}_3^\dagger$. Defining $\mathbf{p}$ as in (5.16), setting $\hat{\mathbf{p}} = \frac{1}{\mu}\mathbf{p}$ and adding a total derivative, we can write the symplectic potential (5.34) associated to the centre of mass motion as

$$\begin{aligned} \langle (\mu J_0 + sP_0), w^{-1}\delta w \rangle - \frac{1}{2}\mu\delta s &= \langle \mathbf{p}, \delta \mathbf{x} \rangle + \langle sP_0, v^{-1}\delta v \rangle = \langle \mathbf{k}, \delta v v^{-1} \rangle - \delta\mu \langle \hat{\mathbf{p}}, \mathbf{x} \rangle \\ &= \langle \mathbf{k}, \delta v v^{-1} \rangle - T\delta\mu, \end{aligned} \quad (7.1)$$

where the vector $\mathbf{k}$ is defined in terms of $\mathbf{x}, \mathbf{p}$ and s as in (3.22) and

$$T = \langle \hat{\mathbf{p}}, \mathbf{x} \rangle. \quad (7.2)$$

From the identification (5.36) it follows that these variables are subject to the identification

$$(T, \mu, \mathbf{k}, v) \sim (T + s, \mu, \mathbf{k}, v e^{-(2\pi - \mu)J_0}). \quad (7.3)$$

Expression (7.1) for the symplectic potential in terms of the variables $T, \mu, v, \mathbf{k}$ implies that the only non-vanishing Poisson brackets of these variables are given by

$$\{k_a, f(v)\} = \frac{d}{d\epsilon} f(e^{-\epsilon J_a} v) \quad \forall f \in \mathcal{C}^\infty(\tilde{L}_3^\uparrow) \quad (7.4)$$

$$\{k_a, k_b\} = \epsilon_{ab}{}^c k_c \quad (7.5)$$

$$\{\mu, T\} = -1. \quad (7.6)$$

The spin of the universe can be expressed in terms of the angular momentum three-vector $\mathbf{k} \in \mathbb{R}^3$ and the Lorentz transformation $v \in \tilde{L}_3^\uparrow$ as

$$s = \langle \mathbf{k}, \text{Ad}(v)J_0 \rangle, \quad (7.7)$$

and it follows from the brackets (7.4), (7.6) that it generates a right-multiplication of the Lorentz element v by a rotation

$$s : \quad v \mapsto v e^{-\epsilon J_0}. \quad (7.8)$$

However, the identification (7.3) implies that there is no value of the flow parameter ϵ for which the flow in phase space generated by the spin is equal to the identity. Instead, we find that the generator of a rotation whose flow by 2π is equal to the identity is the alternative angular momentum

$$J = (1 - \frac{\mu}{2\pi})s, \quad (7.9)$$

which generates the transformation

$$J : \quad (T, \mu, \mathbf{k}, v) \mapsto (T + \epsilon \frac{s}{2\pi}, \mu, \mathbf{k}, v e^{-\epsilon(1 - \frac{\mu}{2\pi})J_0}). \quad (7.10)$$

The phase space function s and J and their relation was first discussed in the metric formalism in a centre of mass frame in [4] and [24].

From expressions (7.4), (7.5) and (7.6) for the Poisson bracket, it is apparent that the Poisson algebra describing the centre of mass degrees of freedom is of a special type, namely the semidirect product of the universal enveloping algebra of the Lie algebra $so(2, 1) \oplus \mathbb{R}$ with the space of $\mathcal{C}^\infty$ -functions on $\tilde{L}_3^\uparrow \times \mathbb{R}$. We considered the quantisation of Poisson algebras of that type in our paper [14]. We can therefore apply our quantisation procedure developed there to construct the quantum algebra and investigate its irreducible Hilbert space representations. From Theorem 3.1 in [14], it follows that the quantum algebra for the Poisson algebra (7.4), (7.5) and (7.6) is the algebra

$$\hat{\mathcal{F}} = U(so(2, 1) \oplus \mathbb{R}) \otimes \mathcal{C}^\infty(\tilde{L}_3^\uparrow \times \mathbb{R}) \quad (7.11)$$

with multiplication

$$\begin{aligned} & ((\mathbf{k}_1, T_1) \otimes F_1) \cdot ((\mathbf{k}_2, T_2) \otimes F_2) \\ &= ((\mathbf{k}_1, T_1) \cdot_U (\mathbf{k}_2, T_2)) \otimes F_1 F_2 + i\hbar (\mathbf{k}_2, T_2) \otimes F_1 \{(\mathbf{k}_1, T_1), F_2\} \\ & \mathbf{k}_1, \mathbf{k}_2 \in so(2, 1), F_1, F_2 \in \mathcal{C}^\infty(\tilde{L}_3^\uparrow \times \mathbb{R}, \mathbb{C}), \end{aligned} \quad (7.12)$$

where the components of the angular momentum three-vector $\mathbf{k}$ and the time T are identified with the generators of the Lie algebra $so(2, 1) \oplus \mathbb{R}$ so that $T_1 = t_1 T$, $T_2 = t_2 T$ with $t_1, t_2 \in \mathbb{R}$. The Lorentz transformation v and mass μ appear as the arguments of the functions $F \in \mathcal{C}^\infty(\tilde{L}_3^\uparrow \times \mathbb{R})$:

$$\{(\mathbf{k}, T), F\}(v, \mu) = \frac{d}{d\epsilon} \Big|_{\epsilon=0} F(e^{-\epsilon k^a J_a} \cdot v, \mu + \epsilon) \quad \forall \mathbf{k} \in so(2, 1), F \in \mathcal{C}^\infty(\tilde{L}_3^\uparrow \times \mathbb{R}). \quad (7.13)$$

As discussed in [14], Sect. 3, the quantum algebra (7.11) is represented on the space

$$V = \mathcal{C}_0^\infty(\tilde{L}_3^\uparrow \times \mathbb{R}), \quad (7.14)$$

of $\mathcal{C}^\infty$ -functions with compact support on the group $\tilde{L}_3^\uparrow \times \mathbb{R}$ according to

$$\Pi(\mathbf{k})\psi(v, \mu) = i\hbar \frac{d}{d\epsilon} \Big|_{\epsilon=0} \psi(e^{-\epsilon k^a J_a} \cdot v, \mu) \quad (7.15)$$

$$\Pi(T)\psi(v, \mu) = i\hbar \frac{\partial}{\partial \mu} \psi(v, \mu)$$

$$\Pi(F)\psi(v, \mu) = F(v, \mu)\psi(v, \mu) \quad \forall \mathbf{k} \in so(2, 1), T \in \mathbb{R}, F \in \mathcal{C}^\infty(\tilde{L}_3^\uparrow \times \mathbb{R}).$$

In particular, the mass operator μ of the universe is identified with the function

$$\mu \in \mathcal{C}^\infty(\tilde{L}_3^\uparrow \times \mathbb{R}) : (v, \tilde{\mu}) \mapsto \tilde{\mu} \quad (7.16)$$

and acts by multiplication

$$\Pi(\mu)\psi(v, \mu) = \mu\psi(v, \mu). \quad (7.17)$$

Furthermore, it follows from (7.7) and (7.15) that the spin operator s acts by right-multiplication of the argument v by an infinitesimal rotation

$$\Pi(s)\psi(v, \mu) = i\hbar \frac{d}{d\epsilon} \Big|_{\epsilon=0} \psi(v e^{-\epsilon J_0}, \mu), \quad (7.18)$$

and, consequently, the action of the angular momentum operator J in (7.9) is given by

$$\Pi(J)\psi(v, \mu) = i\hbar \frac{d}{d\epsilon} \Big|_{\epsilon=0} \psi(v \cdot e^{-\epsilon(1-\frac{\mu}{2\pi})J_0}, \mu). \quad (7.19)$$

To derive a quantisation condition for mass and spin of the universe, we decompose the Hilbert space (7.14) into eigenspaces $V_{\mu s}$ of the (commuting) mass and spin operators:

$$\Pi(s)\psi = s \psi \quad \Pi(\mu)\psi = \mu\psi \quad \forall \psi \in V_{\mu s}. \quad (7.20)$$

From (7.19), it then follows that these spaces are also eigenspaces of the angular momentum operator J . The quantum counterpart of the condition that the flow in phase space generated by the angular momentum J with flow parameter 2π is the requirement that the

unitary operator $e^{2\pi\frac{J}{i\hbar}}$ maps each state to itself. As this operator acts on the eigenspaces $V_{\mu s}$ according to

$$\Pi(e^{2\pi\frac{J}{i\hbar}})\psi = e^{i\frac{(2\pi-\mu)s}{\hbar}}\psi \quad \forall \psi \in V_{\mu s}, \quad (7.21)$$

this implies $(2\pi - \mu)s \in 2\pi\hbar\mathbb{Z}$ or, equivalently, the quantisation condition

$$s = \frac{\hbar n}{1 - \frac{\mu}{2\pi}} \quad \text{with} \quad n \in \mathbb{Z}. \quad (7.22)$$

The condition (7.22) has been found by various authors in more specialised settings. Extrapolating from the study of scattering of a test particle in the background of fixed conical geometry, 't Hooft, Deser, Jackiw and de Sousa Gebert [25, 26, 27] conjectured the quantisation condition (7.22) for the total mass and spin of a two-particle system. Carlip [28] showed that such a condition for the total mass and spin of two spinless particles can be obtained without relying on a semi-classical approximation and is linked to the action of the braid group. This is generalised to spinning particles in [29] where the condition (7.22) is obtained from the condition that the action of the ribbon element of the quantum double $D(\tilde{L}_3^\dagger)$ is identical to a rotation by 2π . The first derivation of (7.22) from first principles is given in the work of Louko and Matschull [30] on the (2+1)-dimensional Kepler problem and the paper [4] on the phase space of a system of n massive, but spinless particles. There it is shown that (7.22) holds for the total mass and spin of, respectively, two and n massive but spinless particles in the metric formulation of (2+1)-dimensional gravity. Our derivation shows that (7.22) holds quite generally for arbitrary genus g and n spinning particles.

Note that in the case of an open universe of genus $g = 0$ with two massive, spinning particles, the discussion in this section amounts to a complete quantisation of the system. According to equation (5.24), the phase space of such a two-particle system is eight-dimensional, and the relative motion of the two particles is determined entirely by the total mass and spin of the system, for details see the discussion in [29]. After the application of the gauge fixing procedure outlined in Sect. 5.4, the phase space is parametrised entirely by the variables $\mu, T, v, \mathbf{k}$ with the identification (7.3), and its Poisson structure is determined by the symplectic potential (7.1). The resulting quantum theory is identical to the one derived in [29] by exploring the analogy of particles in (2+1)-dimensional gravity with anyons.

8 Outlook and conclusion

Based on the treatment of spatial infinity as a distinguished puncture, this paper contains an explicit description of the phase space structure of the Chern-Simons formulation of (2+1)-dimensional gravity in an open universe with vanishing cosmological constant. We restrict attention to universes of product topology $\mathbb{R} \times S$, where S is a two-dimensional oriented manifold, but allow for S to have arbitrary topology and an arbitrary number of punctures. We see generality and explicitness as the main advantages of our approach and expect our results to be useful in the following ways.

The treatment of spatial infinity as a distinguished puncture provides the basis for modelling other physical situations. Here we considered the boundary condition that the universe looks asymptotically like a spinning cone. In that case the holonomy around the distinguished puncture is, in the terminology of Sect. 2, the exponential of an elliptic element of $iso(2, 1)$. However, it is known that the total holonomy for two particles which move very rapidly relative to each other is not of this type [31]. This can be accommodated in our formalism without difficulty. In other contexts it may also be of interest to consider several boundary components, which could be modelled in our formalism by including several distinguished punctures.

Furthermore, we expect our results to be useful in linking different approaches to (2+1)-dimensional gravity, and to provide a unifying viewpoint for existing and future investigations of the phase space structure. This includes studies of closed universes, which can be thought as the special case $\mu = s = 0$ of the open universes considered here (though the discussion of gauge invariance and gauge fixing in Sect. 5 needs to be adapted). In this paper we explained the link between Matschull's and our description of the phase space for n spinless particles on a disc. It would also be interesting to clarify the relation between 't Hooft's polygon approach [32] and the Chern-Simons formulation of (2+1)-dimensional gravity. This should be possible on the basis of our paper since 't Hooft's approach is closely related to Matschull's description of the phase space in [4]. In particular, one should be able to establish an explicit link between our description of the phase space and the coordinates and symplectic structure given in the recent papers [33] and [34], which treat the case where S is a closed surface of arbitrary genus without punctures using the polygon approach.

Finally, we expect the current paper to be useful in investigating the classical dynamics and the quantisation of (2+1)-dimensional gravity. The literature on these topics is vast and we shall make not attempt to survey it here, referring the reader instead to [13] for references. Rigorous and explicit results exist for special cases such as n particle dynamics on a genus $g = 0$ surface or closed universes of the form $\mathbb{R} \times T^2$, where T^2 is the two-torus. There are unifying themes, such as the appearance of the braid group in the analysis of n particle dynamics, but a unified and systematic treatment is still lacking. The current paper provides the starting point for such a treatment since our description of the phase space structure is general, explicit and based on variables which are amenable to a physical interpretation.

Acknowledgements

BJS acknowledges support through an EPSRC advanced fellowship during the early stages of this project and thanks Thomas Strobl for discussions about boundary conditions in field theory and gravity.




Figure 2: A graph on the spatial surface as used in [4]

A Matschull’s description of the phase space in [4]

A.1 General formalism

In this appendix, we summarise the description of phase space and symplectic structure given by Matschull [4] for open universes of genus $g = 0$ with n massive, but spinless particles. We adapt the notation and conventions in [4] to the conventions in the main text. An important difference in conventions is the different sign convention used for the Minkowski metric in [4] (“mostly plus” instead of our “mostly minus”). Hence our pairing $\langle J_a, P_b \rangle = \eta_{ab}$ is minus the trace used in [4]. Also, there is an overall relative sign in the definition of the symplectic structure. Finally, we use the group $\tilde{P}_3^\dagger$ instead of the group $SL(2, \mathbb{R}) \ltimes \mathfrak{sl}_2$ used in [4].

In [4], spatial infinity is incorporated as a connected boundary with an associated boundary condition on dreibein and spin connection. This boundary condition is the requirement that the dreibein and spin connection take the form (3.5) associated to a cone with deficit angle μ and spin s in a region around the boundary. The formalism relies on an ADM-decomposition of the spacetime and an oriented graph Γ embedded into a spatial slice. The embedding is such that every vertex of the graph coincides with one of the particles and each oriented edge either connects two particles or is a spatial half line extending from one particle to the boundary, see Fig. 2. The oriented edges can therefore be grouped into two disjoint sets, the set Γ_+ of oriented *internal* edges that start and end at two vertices of the graph and the set Γ_∞ of *external* edges that start at a vertex and extend to the boundary. For both sets of edges, there are corresponding sets Γ_- and $\Gamma_{-\infty}$ which are obtained by reversing the orientation of each edge in Γ_+ and Γ_∞ , respectively. Among the faces of the graph, one distinguishes *internal polygons*, faces for which all sides are edges connecting particles and *external polygons*, faces for which at least one side is a component of the boundary.

The symplectic structure is described in terms of a symplectic potential defined on an extended phase space of link variables assigned to the oriented graph as follows. To each oriented internal edge $\lambda \in \Gamma_+$, one associates an element $(g_\lambda, \mathbf{z}_\lambda) \in \tilde{P}_3^\dagger$

$$\lambda \in \Gamma_+ \mapsto (g_\lambda, \mathbf{z}_\lambda) \in \tilde{P}_3^\dagger, \quad (\text{A.1})$$

and to the corresponding edge $-\lambda$ with reversed orientation

$$-\lambda \in \Gamma_+ \mapsto (g_\lambda^{-1}, -\text{Ad}(g_\lambda)\mathbf{z}_\lambda) \in \tilde{P}_3^\dagger. \quad (\text{A.2})$$

Furthermore, each oriented external edge $\eta \in \Gamma_\infty$ is equipped with four real parameters M_η, T_η, L_η and φ_η

$$\eta \in \Gamma_\infty \mapsto (M_\eta, T_\eta, L_\eta, \varphi_\eta). \quad (\text{A.3})$$

In terms of these link variables the Lorentz component of the holonomy around a particle π with respect to a base point in an adjacent face Δ is given by

$$u_{\pi, \Delta} = \prod_{\lambda \in \Delta \cap \pi} g_\lambda, \quad (\text{A.4})$$

where $\lambda \in \Delta \cap \pi$ denotes the set of all edges that either start or end at the vertex associated to the particle π and are a side of the polygon Δ . The total mass μ and total spin s of the universe can be expressed in terms of the link variables as

$$\mu = \sum_{\eta \in \Gamma_\infty} M_\eta \quad (\text{A.5})$$

$$s = - \sum_{\lambda \in \Gamma_+} \langle \mathbf{z}_\lambda - \text{Ad}(g_\lambda)\mathbf{z}_\lambda, J_0 \rangle. \quad (\text{A.6})$$

The symplectic structure on the extended phase space is given as the exterior derivative of the symplectic potential

$$-\Theta_m(g_\lambda, \mathbf{z}_\lambda, M_\eta, T_\eta, L_\eta, \varphi_\eta) = - \sum_{\eta \in \Gamma_\infty} (T_\eta \delta M_\eta + L_\eta \delta \varphi_\eta) - \sum_{\lambda \in \Gamma_+} \langle g_\lambda^{-1} \delta g_\lambda, \mathbf{z}_\lambda \rangle, \quad (\text{A.7})$$

where we have inserted an overall minus sign in the result given in [4] and taken into account the different sign convention for the pairing $\langle, \rangle$ to make the formula (A.7) compatible with the conventions used in this paper. The physical phase space and its symplectic structure are obtained from the extended phase space by imposing four sets of first-class constraints and by dividing by the associated gauge transformations. The first set of constraints states that for each internal polygon the sum of the relative position vectors $\mathbf{z}_\lambda$ associated to its sides must vanish

$$\mathbf{Z}_\Delta = \sum_{\lambda \in \Delta} \mathbf{z}_\lambda \approx 0 \quad \forall \Delta \in \Pi_0. \quad (\text{A.8})$$

The associated gauge transformations are Lorentz transformations $v \in \tilde{L}_3^\uparrow$ acting on the edges of the polygon via Δ

$$\begin{aligned} \mathbf{Z}_\Delta : \quad g_\lambda &\mapsto g_\lambda v^{-1} && \text{for } \lambda \in \Delta \\ g_\lambda &\mapsto v g_\lambda && \text{for } -\lambda \in \Delta \\ \mathbf{z}_\lambda &\mapsto \text{Ad}(v) \mathbf{z}_\lambda && \text{for } \lambda \in \Delta \text{ or } -\lambda \in \Delta. \end{aligned} \quad (\text{A.9})$$

The second set of constraints refers to external polygons Δ

$$\mathcal{Z}_\Delta = T_{\eta'} - T_\eta + \frac{s}{2\pi}(\varphi_{\eta'} - \varphi_\eta) - \sum_{\lambda \in \Delta \cap (\Gamma_+ \cup \Gamma_-)} \langle \mathbf{z}_\lambda, J_0 \rangle \approx 0 \quad \forall \Delta \in \Pi_\infty, \quad (\text{A.10})$$

where η' denotes the edge consecutive to edge η in counterclockwise direction, see Fig. 2, and $\Delta \cap (\Gamma_+ \cup \Gamma_-)$ the sides of the external polygon that correspond to internal edges. It generates gauge transformations that shift the parameters M_η and L_η of the external edges in the polygon Δ as well as a rotating all internal polygons by a real parameter τ

$$\begin{aligned} \mathcal{Z}_\Delta : \quad M_{\eta'} &\mapsto M_{\eta'} - \tau, \quad M_\eta \mapsto M_\eta + \tau \\ L_{\eta'} &\mapsto L_{\eta'} + \frac{s}{2\pi}\tau, \quad L_\eta \mapsto L_\eta - \frac{s}{2\pi}\tau \\ g_\lambda &\mapsto e^{-\frac{\tau}{2\pi}(\varphi_{\eta'} - \varphi_\eta)J_0} g_\lambda e^{\frac{\tau}{2\pi}(\varphi_{\eta'} - \varphi_\eta)J_0}, \quad \mathbf{z}_\lambda \mapsto \text{Ad}(e^{-\frac{\tau}{2\pi}(\varphi_{\eta'} - \varphi_\eta)J_0}) \mathbf{z}_\lambda \text{ for } \lambda, -\lambda \notin \Delta \\ g_\lambda &\mapsto e^{-\frac{\tau}{2\pi}(\varphi_{\eta'} - \varphi_\eta)J_0} g_\lambda e^{(\frac{\tau}{2\pi}(\varphi_{\eta'} - \varphi_\eta) - \tau)J_0}, \quad \mathbf{z}_\lambda \mapsto \text{Ad}(e^{(-\frac{\tau}{2\pi}(\varphi_{\eta'} - \varphi_\eta) + \tau)J_0}) \mathbf{z}_\lambda \\ &\quad \text{for } \lambda \in \Delta \cap (\Gamma_+ \cup \Gamma_-). \end{aligned} \quad (\text{A.11})$$

Furthermore, there is an additional constraint for each external edge $\eta \in \gamma_\infty$, which relates the associated parameters M_η, L_η to the total spin of the asymptotic cone

$$\mathcal{J}_\eta = L_\eta + \frac{s}{2\pi}M_\eta \approx 0 \quad \forall \eta \in \Gamma_\infty. \quad (\text{A.12})$$

The gauge transformations for these constraints shift the variables T_η and φ_η of this edge and, again, also rotate all internal polygons

$$\begin{aligned} \mathcal{J}_\eta : \quad T_\eta &\mapsto T_\eta + \frac{s}{2\pi}\psi, \quad \varphi_\eta \mapsto \varphi_\eta - \psi \\ g_\lambda &\mapsto e^{-\frac{M_\eta}{2\pi}\psi J_0} g_\lambda e^{\frac{M_\eta}{2\pi}\psi J_0}, \quad \mathbf{z}_\lambda \mapsto \text{Ad}(e^{-\frac{M_\eta}{2\pi}\psi J_0}) \mathbf{z}_\lambda \quad \forall \lambda \in \Gamma_+ \cup \Gamma_-, \end{aligned} \quad (\text{A.13})$$

where $\psi \in \mathbb{R}$. Finally, there is set of mass constraints for each particle. Parametrising the holonomy (A.4) around the particle in terms of a momentum three vector $\mathbf{p}_{\pi,\Delta}$, these mass constraints read

$$C_{\pi,\Delta} = \sqrt{\mathbf{p}_{\pi,\Delta}^2} - \mu_\pi \approx 0 \quad \text{where} \quad u_{\pi,\Delta} = e^{-p_{\pi,\Delta}^a J_a}. \quad (\text{A.14})$$

The associated gauge transformations translate the vectors $\mathbf{z}_\lambda$ of internal edges λ starting or ending at the particle and shift the parameter T_η of the external edges starting at the particle

$$\begin{aligned} C_{\pi,\Delta} : \quad \mathbf{z}_\lambda &\mapsto \mathbf{z}_\lambda - \tau \hat{\mathbf{p}}_{\pi,\Delta} && \text{if } \lambda \in \Delta \text{ and points to vertex } \pi \\ \mathbf{z}_\lambda &\mapsto \mathbf{z}_\lambda + \tau \hat{\mathbf{p}}_{\pi,\Delta} && \text{if } \lambda \in \Delta \text{ and starts at vertex } \pi \\ T_\eta &\mapsto T_\eta + \tau \langle \hat{\mathbf{p}}_{\pi,\Delta}, J_0 \rangle && \text{if } \eta \in \Delta \text{ and starts at vertex } \pi, \quad \tau \in \mathbb{R}, \end{aligned} \quad (\text{A.15})$$

The symplectic potential (A.7) on the extended phase space becomes

$$-\Theta_m(g_i, \mathbf{z}_i, \mu, s, T_0, \varphi_0) = -T_0 \delta \mu - L_0 \delta \varphi_0 - \sum_{i=1}^{n-1} \langle g_i^{-1} \delta g_i, \mathbf{z}_i \rangle. \quad (\text{A.20})$$

To obtain the physical phase space, one has to impose the constraints (A.8), (A.10), (A.12) and (A.14) and to divide by the associated gauge transformations. As the graph in Fig. 3 has only a single face and one external edge, the constraints (A.8) and (A.10) are satisfied trivially. However, there is an external edge and therefore an associated constraint of the form (A.12). Taking into account (A.18), this constraint becomes

$$\mathcal{J}_0 = L_0 + \frac{\mu s}{2\pi} \approx 0, \quad (\text{A.21})$$

where s is the spin of the universe given as a function of the variables $g_i, \mathbf{z}_i$ by (A.19). The associated gauge transformation (A.13) is precisely the transformation (6.18) given in the main text.

The remaining constraints are the n mass constraints (A.14) the particles. With the parametrisation $\tilde{u}_{M_i} = e^{-\tilde{p}_{M_i}^a J_a}$ for the Lorentz holonomies (A.17) around each particle, they read

$$C_i : \quad \sqrt{\tilde{\mathbf{p}}_{M_i}^2} - \mu_i \approx 0 \quad i = 1, \dots, n, \quad (\text{A.22})$$

and the gauge transformations (A.15) now take the form (6.19) given in the main text.

The physical phase space is obtained from the extended phase space, parametrised by the $n - 1$ Poincaré elements $(g_i, \mathbf{z}_i)$ and the parameters $M_0 = \mu, T_0, L_0$ and φ_0 and carrying symplectic potential (A.20), by imposing the n mass constraints (A.22) for each particle and the constraint (A.21) and dividing by the associated gauge transformations (6.19) and (6.18). It therefore has the dimension $6(n - 1) + 4 - 2(n + 1) = 4n - 4$.

References

- [1] A. Achúcarro, P. Townsend, A Chern–Simons action for three-dimensional anti-de Sitter supergravity theories, *Phys. Lett. B* 180 (1986) 85–100.
- [2] E. Witten, 2+1 dimensional gravity as an exactly soluble system, *Nucl. Phys. B* 311 (1988) 46–78.
- [3] S. Deser, R. Jackiw, G. 't Hooft, Three-dimensional Einstein gravity: dynamics of flat space, *Ann. Phys.* 152 (1984) 220–35.
- [4] H. -J. Matschull, The phase space structure of multiparticle models in 2+1 gravity, *Class. Quant. Grav.* 18 (2001) 3497–3560.
- [5] S. Carlip, Statistical mechanics of the (2+1)-dimensional black hole, *Phys. Rev. D* 51 (1995) 632–637.

- [6] M. Banados and A. Gomberoff, Black hole entropy in the Chern-Simons formulation of (2+1) gravity Phys. Rev. D55 (1997) 6162–6167.
- [7] E. Witten, Quantum Field Theory and the Jones Polynomial, Commun. Math. Phys. 121 (1989) 351–399.
- [8] P. de Sousa Gerbert, On spin and (quantum) gravity in 2+1 dimensions, Nucl. Phys. B346 (1990) 440–472.
- [9] C. Meusburger, B. J. Schroers, Phase space structure of Chern-Simons theory with a non-standard puncture, EMPG-05-07, hep-th/0505143.
- [10] A. Y. Alekseev and A. Z. Malkin, Symplectic structure of the moduli space of flat connections on a Riemann surface, Commun. Math. Phys. 169 (1995) 99–119.
- [11] C. Meusburger, B. J. Schroers, Poisson structure and symmetry in the Chern-Simons formulation of (2+1)-dimensional gravity, Class. Quant. Grav. 20 (2003) 2193–2233.
- [12] V. V. Fock, A. A. Rosly, Poisson structures on moduli of flat connections on Riemann surfaces and r -matrices, *ITEP preprint* (1992) 72-92 (see also math.QA/9802054).
- [13] S. Carlip, Quantum gravity in 2+1 dimensions, Cambridge University Press, Cambridge, 1998.
- [14] C. Meusburger, B. J. Schroers, The quantisation of Poisson structures arising in Chern-Simons theory with gauge group $G \ltimes \mathfrak{g}^*$, Adv. Theor. Math. Phys. 7 (2004) 1003–1043.
- [15] C. Meusburger, B. J. Schroers, Mapping class group actions in Chern-Simons theory with gauge group $G \ltimes \mathfrak{g}^*$, Nucl. Phys. B 706 (2005) 569–597.
- [16] A. W. Knap, Lie groups Beyond an introduction, Birkhäuser, Boston, 1996.
- [17] H. -J. Matschull, On the relation between (2+1) Einstein gravity and Chern-Simons Theory, Class. Quant. Grav. 16 (1999) 2599–609.
- [18] P. Schaller and T. Strobl, Diffeomorphisms versus non abelian gauge transformations: an example of (1+1)-dimensional gravity, Phys. Lett. B337 (1994) 266–270.
- [19] W. P. Thurston, Three-dimensional geometry and topology, Princeton University Press, Princeton, 1997.
- [20] R. W. Sharpe, Differential Geometry, Springer Verlag, New York, 1996.
- [21] D. Giulini, C. Kiefer, H. D. Zeh, Symmetries, superselection rules and decoherence, Phys. Lett. A 199 (1995) 291.
- [22] T. Regge, C. Teitelboim, Role of surface integrals in the Hamiltonian formulation of general relativity, Ann. Phys. 88 (1974) 286–318.

- [23] D. Giulini, Asymptotic symmetry groups of long-ranged gauge configurations, *Mod. Phys. Lett. A* 10 (1995) 2059–2070.
- [24] J. Louko and H. -J. Matschull, (2+1)-dimensional Einstein-Kepler problem in the centre-of-mass frame *Class. Quant. Grav.* 17 (2000) 1847–1873.
- [25] G. 't Hooft, Non-perturbative 2 particle scattering amplitude in 2+1 dimensional quantum gravity, *Commun. Math. Phys.* 117 (1988) 685–700.
- [26] S. Deser, R. Jackiw, Classical and quantum scattering on a cone, *Commun. Math. Phys.* 118 (1988) 495–509.
- [27] P. de Sousa Gerbert, R. Jackiw, Classical and quantum scattering on a spinning cone, *Commun. Math. Phys.* 124 (1989) 229–260.
- [28] S. Carlip, Exact quantum scattering in (2+1)-dimensional gravity, *Nucl. Phys. B* 324 (1989) 106–22.
- [29] F. A. Bais, N. M. Muller, B. J. Schroers, Quantum group symmetry and particle scattering in (2+1)-dimensional quantum gravity, *Nucl. Phys. B* 640 (2002) 3–45.
- [30] J. Louko and H. J. Matschull, The 2+1 Kepler problems and its quantisation, *Class. Quant. Grav.* 18 (2001) 2731–2784.
- [31] J. R. Gott, Closed timelike curves produced by pairs of moving cosmic strings: exact solutions, *Phys. Rev. Lett.* 66 (1991) 1126–9.
- [32] G. 't Hooft, Causality in (2+1)-dimensional gravity, *Class. Quant. Grav.* 9, (1992) 1335–1348.
- [33] Z. Kadar and R. Loll, (2+1) gravity for higher genus in the polygon model, *Class. Quant. Grav.* 21 (2004) 2465–2491.
- [34] Z. Kadar, Polygon model form first order gravity, *Class. Quant. Grav.* 22 (2005) 809–824.